



NIGHTTIME DOSING



DAYTIME COVERAGE

NEW
ADDITIONAL
INDICATION

**GOCOVRI® IS
NOW FDA APPROVED
TO ALSO TREAT 'OFF'
EPISODES IN PEOPLE
TAKING LEVODOPA**

Not an actual patient.

If you're living with Parkinson's disease
and experience motor complications

**GOCOVRI® COULD MEAN THE
DIFFERENCE BETWEEN
GETTING UP AND
GETTING OUT**

ONCE DAILY AT BEDTIME
GOCOVRI®
(amantadine) extended release capsules
68.5 mg | 137 mg

WHAT IS GOCOVRI®?

GOCOVRI is a prescription medicine used:

- for the treatment of dyskinesia (sudden uncontrolled movements) in people with Parkinson's disease who are treated with levodopa therapy or levodopa therapy with other medicines that increase the effects of dopamine in the brain.
- with levodopa and carbidopa in people with Parkinson's disease who are having "off" episodes.

It is not known if GOCOVRI is safe and effective in children.

IMPORTANT SAFETY INFORMATION

DO NOT take GOCOVRI if you have severe kidney problems.

WHAT SHOULD I AVOID WHILE TAKING GOCOVRI®?

Do not stop or change the dose of GOCOVRI before talking with your doctor. Call your healthcare provider if you have symptoms of withdrawal such as fever, confusion, or severe muscle stiffness.

Do not drink alcohol while taking GOCOVRI. It can increase your chances of getting serious side effects.


DO NOT drive, operate machinery, or do other dangerous activities until you know how GOCOVRI affects you.

If you take too much GOCOVRI, call your doctor or go to the nearest hospital emergency room right away.

**Please see Important Safety Information throughout
and full Patient Prescribing Information on pages 15-19.**

GOOD ON time can be limited by OFF time and dyskinesia

Some important terms to know about movement control in PD



IMPORTANT SAFETY INFORMATION (continued) WHAT SHOULD I AVOID WHILE TAKING GOCOVRI? (continued)

Do not drive, operate machinery, or do other dangerous activities until you know how GOCOVRI affects you.

If you take too much GOCOVRI, call your doctor or go to the nearest hospital emergency room right away.

WHAT ARE THE POSSIBLE SIDE EFFECTS OF GOCOVRI?

- **Falling asleep during normal activities.** Activities may include driving, talking, or eating. You may fall asleep without being drowsy or warning.

IMPORTANT SAFETY INFORMATION (continued) WHAT ARE THE POSSIBLE SIDE EFFECTS OF GOCOVRI? (continued)

- **Suicidal thoughts or actions and depression.** Tell your doctor if you have new or sudden changes in mood, behaviors, thoughts, or feelings, including thoughts about hurting yourself or ending your life.

Please see Important Safety Information throughout and full Patient Prescribing Information on pages 15-19.

Dyskinesia may have an effect on daily routines and activities

Studies showed up to 73% of people with Parkinson's who have dyskinesia said it had a functional impact on them^{1*}



WALKING



HOBBIES



HANDWRITING



DRESSING



HYGIENE



EATING

Dyskinesia can also have a social and emotional impact

Studies showed 72% of PWP with dyskinesia said it had a social impact on them^{1*}

*Impact of dyskinesia on activities of daily living in Parkinson's disease: Results from pooled phase 3 ADS-5102 clinical trials.

In a 2012 survey, 40 respondents identified their main concerns²:

- Embarrassment
- Avoiding family events and gatherings
- Family tension

IMPORTANT SAFETY INFORMATION (continued)
WHAT ARE THE POSSIBLE SIDE EFFECTS OF GOCOVRI?
(continued)

- **Hallucinations.** GOCOVRI can cause or worsen hallucinations (seeing or hearing things that are not real) or psychotic behavior.

Please see Important Safety Information throughout and full Patient Prescribing Information on pages 15-19.



"Dyskinesia and the loss of control of my movement affected my ability to play musical instruments. GOCOVRI® has helped me control my dyskinesia, which has given me a chance to start again."


Kendall, real patient

GOCOVRI® increased GOOD ON time by reducing dyskinesia and OFF time

GOCOVRI® clinical studies measured results in 2 ways

Based on UDysRS score vs placebo:

- The Unified Dyskinesia Rating Scale (UDysRS) scores were used to measure the severity of dyskinesia
- It looked at how dyskinesia impacted daily activities



IMPORTANT SAFETY INFORMATION (continued) WHAT ARE THE POSSIBLE SIDE EFFECTS OF GOCOVRI? (continued)

- **Feeling dizzy, faint or lightheaded, especially when you stand up (orthostatic hypotension).** Lightheadedness or fainting may happen when getting up too quickly after long periods of time, when first starting GOCOVRI, or if your dose has been increased.
- **Unusual urges.** Examples include gambling, sexual urges, spending money, binge eating, and the inability to control them.

Please see Important Safety Information throughout and full Patient Prescribing Information on pages 15-19.

Based on patient diary data vs placebo:

The patient diaries were used to track movement throughout the day

SAMPLE DIARY PLOT FOR A SINGLE PATIENT AT STUDY BASELINE THROUGHOUT THE WAKING DAY



STUDY RESULTS: PATIENT DIARY DATA VS PLACEBO



Take GOCOVRI® at bedtime for dyskinesia control throughout the day



You take
GOCOVRI®
at bedtime



While you sleep,
GOCOVRI® releases
slowly in your body



By the time you wake up in the
morning, GOCOVRI® has reached
high levels for dyskinesia
control throughout the day



GOCOVRI® is added to your existing levodopa therapy

GOCOVRI® is designed to help!

It's amantadine—but in a different formulation that is specifically indicated to treat your dyskinesia.

It's different because you take it at bedtime and there is a slow lag in release, so that you reach high levels when you wake, before your levodopa dose. GOCOVRI® slowly releases throughout the day before levels taper off in the evening.

IMPORTANT SAFETY INFORMATION (continued) **WHAT ARE THE POSSIBLE SIDE EFFECTS OF GOCOVRI?** (continued)

The most common side effects of GOCOVRI include dry mouth, swelling of legs and feet, constipation, and falls. If you or your family notices that you are developing any new, unusual or sudden changes in behavior, related symptoms, or side effects, tell your healthcare provider right away.

These are not all the possible side effects of GOCOVRI. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Please see Important Safety Information throughout and full Patient Prescribing Information on pages 15-19.



**"I take my GOCOVRI® in the evening
at bedtime, and it lasts throughout
the next day."**

Ervin, real patient

Our 3-step process to getting GOCOVRI®

STEP **1** **A CALL FROM THE PHARMACY**
If your doctor determines GOCOVRI® is right for you, GOCOVRI Onboard® can help. To receive GOCOVRI®, our specialty pharmacy partner needs to speak to you

STEP **2** **GOCOVRI® IS EXPRESS-MAILED TO YOU**
Once our specialty pharmacy confirms your address over the phone, they will arrange express delivery of GOCOVRI® directly to your door, with no need to go to the pharmacy

STEP **3** **FOLLOW-UP CALL**
After you receive GOCOVRI®, your GOCOVRI® Care Coordinator will call you to answer any questions you may have and to discuss next steps

IMPORTANT SAFETY INFORMATION (continued) **TELL YOUR DOCTOR ABOUT ALL OF YOUR MEDICAL CONDITIONS, INCLUDING IF YOU:**

- have kidney problems.
- have unexpected or unpredictable sleepiness, sleep disorders, or currently take medication to help you sleep or makes you drowsy.
- are pregnant or plan to become pregnant or are breastfeeding or plan to breastfeed. GOCOVRI may harm your unborn baby and can pass into your breastmilk.



**Call 1-844-GOCOVRI
(1-844-462-6874)
and press 1
for a GOCOVRI® Care
Coordinator**

Monday-Friday, 8 AM-8 PM ET

**Please see Important Safety Information throughout
and full Patient Prescribing Information on pages 15-19.**

GOCOVRI Onboard® may be able to help you get access to GOCOVRI®



**GOCOVRI
ONBOARD®**
support focused on you

If you and your doctor determine that GOCOVRI® is right for you, GOCOVRI Onboard® can help with accessing our patient support programs.

Your GOCOVRI® Care Coordinator will work with you to see if you are eligible for assistance with paying for GOCOVRI®.*

*This offer is not valid for prescriptions that are eligible to be reimbursed, in whole or in part, by Medicaid, Medicare, TRICARE, the VA healthcare.

IMPORTANT SAFETY INFORMATION (continued) **TELL YOUR DOCTOR ABOUT ALL OF YOUR MEDICAL CONDITIONS, INCLUDING IF YOU:** (continued)

Tell your doctor about all the medicines you take.

Include prescription and over-the-counter medicines, vitamins, and herbal supplements.

Especially tell your doctor if you take medicines like sodium bicarbonate, or have had or are planning to have a live flu vaccination (nasal spray). You can receive the flu vaccination shot but should not get a live flu vaccine while taking GOCOVRI.

We're here for you!



\$20 CO-PAY PROGRAM

If you have a commercial health plan and participate in the GOCOVRI® Co-pay Assistance Program, you'll pay \$20 per prescription



PATIENT ASSISTANCE PROGRAM

If you don't have insurance, your insurance doesn't cover* GOCOVRI®, or if you can't afford GOCOVRI®, the Patient Assistance Program may help you get GOCOVRI® at no cost

INDEPENDENT CHARITABLE FOUNDATIONS PROGRAM

If you have government-sponsored insurance, independent organizations and/or other resources may be available to help



**Call 1-844-GOCOVRI
(1-844-462-6874)
and press 1**
for a GOCOVRI® Care
Coordinator
Monday-Friday, 8 AM-8 PM ET

Please see Important Safety Information throughout and full Patient Prescribing Information on pages 15-19.

Full Patient Prescribing Information



“Since taking GOCOVRI®, I’ve experienced a reduction in my dyskinesic episodes allowing my wife, Gayle, and me to better attend church, visit our family, and volunteer, which we love.”

Ervin, real patient

PATIENT INFORMATION

GOCOVRI® (goh-KUV-ree)

(amantadine) extended release capsules, for oral use

What is GOCOVRI?

GOCOVRI is a prescription medicine used:

- for the treatment of dyskinesia (sudden uncontrolled movements) in people with Parkinson's disease who are treated with levodopa therapy or levodopa therapy with other medicines that increase the effects of dopamine in the brain.
- with levodopa and carbidopa in people with Parkinson's disease who are having “off” episodes.

It is not known if GOCOVRI is safe and effective in children.

Do not take GOCOVRI if you have severe kidney problems

Before you take GOCOVRI, tell your doctor about all of your medical conditions, including if you:

- have kidney problems.
- have daytime sleepiness from a sleep disorder, have unexpected or unpredictable sleepiness or periods of sleep, take a medicine to help you sleep, or take any medicine that makes you drowsy.
- have mental problems, such as suicidal thoughts, depression, or hallucinations.
- have unusual urges including gambling, increased sex drive, compulsive eating, or compulsive shopping.
- drink alcoholic beverages. This may increase your chances of becoming drowsy or sleepy while taking GOCOVRI.
- are pregnant or plan to become pregnant. GOCOVRI may harm your unborn baby.

Full Patient Prescribing Information

(continued)

- are breastfeeding or plan to breastfeed. GOCOVRI can pass into your breastmilk. Talk to your doctor about the best way to feed your baby if you take GOCOVRI.

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Especially tell your doctor if you:

- take medicines like sodium bicarbonate.
- have had or are planning to have a live flu (influenza) vaccination (nasal spray). You can receive the influenza vaccination shot but should not get a live influenza vaccine while taking GOCOVRI.

Know the medicines you take. Keep a list of them to show your doctor and pharmacist when you get a new medicine.

How should I take GOCOVRI?

- Take GOCOVRI as exactly as your doctor tells you to.
- Start GOCOVRI with 1 capsule at bedtime. Your doctor may change your dose if needed. If your doctor tells you to take 2 capsules, take them together at bedtime.
- **Do not** stop or change GOCOVRI before talking with your doctor. Call your doctor if you have symptoms of withdrawal such as fever, confusion, or severe muscle stiffness.
- GOCOVRI may be taken with food or without food.
- Swallow GOCOVRI capsules whole. **Do not** crush, chew, or divide.
- GOCOVRI capsules may be opened and sprinkled on applesauce before swallowing. Sprinkle all of the medicine in the capsule on the applesauce. Take all of the medicine right away without chewing.

- If you miss a dose of GOCOVRI, do not take an extra dose. Take your usual dose of GOCOVRI on the next day at bedtime.
- If you have forgotten to take GOCOVRI for several days, contact your doctor.
- **Do not** drink alcohol with your dose of GOCOVRI.
- **Do not** use GOCOVRI capsules that are damaged or show signs of tampering.
- If you take too much GOCOVRI, call your doctor or go to the nearest hospital emergency room right away.

What should I avoid while taking GOCOVRI?

- **Do not** drive, operate machinery, or do other dangerous activities until you know how GOCOVRI affects you.
- **Do not** drink alcohol while taking GOCOVRI. It can increase your chances of getting serious side effects.

What are the possible side effects of GOCOVRI?

GOCOVRI may cause serious side effects, including:

- **falling asleep during normal activities.** You may fall asleep while doing normal activities such as driving a car, talking, or eating while taking GOCOVRI or other medicines that treat Parkinson's disease. You may fall asleep without being drowsy or without warning. This may result in having accidents. Your chances of falling asleep while doing normal activities while taking GOCOVRI are greater if you take other medicines that cause drowsiness. Tell your doctor right away if this happens.
- **suicidal thoughts or actions and depression.** Some people taking GOCOVRI have had suicidal thoughts, attempted suicide, or depression. Tell your doctor if you have new or sudden changes in mood, behaviors, thoughts, or feelings, including thoughts about hurting yourself or ending your life.

Full Patient Prescribing Information

(continued)

- **hallucinations.** GOCOVRI can cause or worsen hallucinations (seeing or hearing things that are not real) or psychotic behavior. Contact your doctor if you have hallucinations.
- **feeling dizzy, faint or lightheaded, especially when you stand up (orthostatic hypotension).** Lightheadedness or fainting may happen when getting up too quickly from a sitting or lying position especially after long periods of time, when first starting GOCOVRI, or if your dose has been increased. Contact your doctor if you become lightheaded or faint when standing up.
- **unusual urges.** Some people taking GOCOVRI get urges to behave in a way unusual for them. Examples of this are an unusual urge to gamble, increased sexual urges, strong urges to spend money, binge eating and the inability to control these urges. If you notice or your family notices that you are developing any unusual behaviors, talk to your doctor. The most common side effects of GOCOVRI include dry mouth, swelling of legs and feet, constipation, and falls.

These are not all the possible side effects of GOCOVRI. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store GOCOVRI?

- Store GOCOVRI at room temperature between 68°F to 77°F (20°C to 25°C).
- Keep GOCOVRI out of the reach of children.

General information about the safe and effective use of GOCOVRI.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use GOCOVRI for a condition for which it was not prescribed. Do not give GOCOVRI to other people, even if they have the same symptoms that you have. It may harm them. You can ask your pharmacist or doctor for information about GOCOVRI that is written for health professionals.

What are the ingredients in GOCOVRI?

Active ingredient: amantadine hydrochloride

Inactive ingredients: copovidone, ethylcellulose, hypromellose, magnesium stearate, medium chain triglycerides, microcrystalline cellulose, povidone, and talc

Manufactured for:
Adamas Pharma LLC. Emeryville, CA 94608

For more information, go to
www.GOCOVRI.com or call 1-833-223-2627.

This Patient Information has been approved
by the U.S. Food and Drug Administration
Issued: 3/2020

Questions to consider asking your doctor at your next appointment

- I sometimes experience involuntary, erratic, writhing movements of the face, arms, legs, or trunk. Could this be dyskinesia?
- I have heard that one way to manage dyskinesia is to adjust the levodopa dose. Will adjusting the levodopa dose cause more OFF time? What else could be done?
- Dyskinesia/OFF time is interfering with my daily activities. What are my treatment options, and could GOCOVRI[®] be right for me?
- What are the side effects of GOCOVRI[®]?

Create your own customized discussion guide for your visit with your doctor and find other resources at **GOCOVRI.com**

Please see full Patient Prescribing Information on pages 15–19.

References: **1.** Pahwa R, Isaacson S, Jimenez-Shaheed J, et al. Impact of dyskinesia on activities of daily living in Parkinson's disease: Results from pooled phase 3 ADS-5102 clinical trials. *Parkinsonism Relat Disord.* 2019;60:118–125. **2.** Data on file. Adamas Pharma LLC, Emeryville, CA.



Gocovri, the Gocovri logo, Gocovri Onboard, the Gocovri Onboard logo, Adamas, and the Adamas logo are trademarks of Adamas Pharmaceuticals, Inc. or its related companies. © 2021 Adamas Pharmaceuticals, Inc. or its related companies. All rights reserved. GOC-0729 V2 2/21