



TRODELVY™

sacituzumab govitecan-hziy

180 mg for injection

**FULL PRESCRIBING  
INFORMATION**

## HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use TRODELVY™ safely and effectively. See full prescribing information for TRODELVY™

TRODELVY™ (sacituzumab govitecan-hziy) for injection, for intravenous use

Initial U.S. Approval: 2020

### WARNING: NEUTROPENIA AND DIARRHEA

- Severe neutropenia may occur. Withhold TRODELVY for absolute neutrophil count below 1500/mm<sup>3</sup> or neutropenic fever. Monitor blood cell counts periodically during treatment. Consider G-CSF for secondary prophylaxis. Initiate anti-infective treatment in patients with febrile neutropenia without delay. (5.1)
- Severe diarrhea may occur. Monitor patients with diarrhea and give fluid and electrolytes as needed. Administer atropine, if not contraindicated, for early diarrhea of any severity. At the onset of late diarrhea, evaluate for infectious causes and, if negative, promptly initiate loperamide. If severe diarrhea occurs, withhold TRODELVY until resolved to ≤ Grade 1 and reduce subsequent doses. (2.3, 5.2)

### INDICATIONS AND USAGE

TRODELVY is a Trop-2-directed antibody and topoisomerase inhibitor conjugate indicated for the treatment of adult patients with metastatic triple-negative breast cancer (mTNBC) who have received at least two prior therapies for metastatic disease. (1)

This indication is approved under accelerated approval based on tumor response rate and duration of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in confirmatory trials. (1,14)

### DOSAGE AND ADMINISTRATION

- Do NOT substitute TRODELVY for or use with other drugs containing irinotecan or its active metabolite SN-38. (2.1)
- For intravenous infusion only. Do not administer as an intravenous push or bolus.
- The recommended dose is 10 mg/kg once weekly on Days 1 and 8 of continuous 21-day treatment cycles until disease progression or unacceptable toxicity. (2.2)
- Premedication for prevention of infusion reactions and prevention of chemotherapy-induced nausea and vomiting is recommended. (2.2)
- Monitor patients during the infusion and for at least 30 minutes after completion of infusion. Treatment interruption and/or dose reduction may be needed to manage adverse reactions. (2.2)
- See Full Prescribing Information for preparation and administration instructions. (2.4)

### DOSAGE FORMS AND STRENGTHS

For injection: 180 mg lyophilized powder in single-dose vials for reconstitution. (3)

### CONTRAINDICATIONS

Severe hypersensitivity reaction to TRODELVY. (4, 5.3)

### WARNINGS AND PRECAUTIONS

- Hypersensitivity: Hypersensitivity reactions including severe anaphylactic reactions have been observed. Monitor patients for infusion-related reactions. Permanently discontinue TRODELVY if severe or life-threatening reactions occur. (5.3)
- Nausea/Vomiting: Use antiemetic preventive treatment and withhold TRODELVY for patients with Grade 3 nausea or Grade 3-4 vomiting at the time of scheduled treatment. (5.4)
- Patients with Reduced UGT1A1 Activity: Individuals who are homozygous for the uridine diphosphate-glucuronosyl transferase 1A1 (UGT1A1)\*28 allele are at increased risk for neutropenia following initiation of TRODELVY treatment. (5.5)
- Embryo-Fetal Toxicity: TRODELVY can cause fetal harm. Advise patients of potential risk to a fetus and to use effective contraception. (5.6, 8.1, 8.3)

### ADVERSE REACTIONS

Most common adverse reactions (incidence ≥25%) in patients with mTNBC are nausea, neutropenia, diarrhea, fatigue, anemia, vomiting, alopecia, constipation, rash, decreased appetite, and abdominal pain. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Immunomedics, Inc. at 1-888-983-4668 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

### DRUG INTERACTIONS

- UGT1A1 inhibitors or inducers: Avoid concomitant use. (7)

### USE IN SPECIFIC POPULATIONS

- Lactation: Advise not to breastfeed. (8.2)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 04/2020

## FULL PRESCRIBING INFORMATION: CONTENTS\*

WARNING: NEUTROPENIA AND DIARRHEA	8	USE IN SPECIFIC POPULATIONS
1 INDICATIONS AND USAGE		8.1 Pregnancy
		8.2 Lactation
2 DOSAGE AND ADMINISTRATION		8.3 Females and Males of Reproductive Potential
2.1 Important Use Information		8.4 Pediatric Use
2.2 Recommended Dose and Schedule		8.5 Geriatric Use
2.3 Dose Modifications for Adverse Reactions		8.6 Hepatic Impairment
2.4 Preparation for Administration		
3 DOSAGE FORMS AND STRENGTHS		10 OVERDOSAGE
4 CONTRAINDICATIONS		11 DESCRIPTION
5 WARNINGS AND PRECAUTIONS		12 CLINICAL PHARMACOLOGY
5.1 Neutropenia		12.1 Mechanism of Action
5.2 Diarrhea		12.2 Pharmacodynamics
5.3 Hypersensitivity		12.3 Pharmacokinetics
5.4 Nausea and Vomiting		12.5 Pharmacogenomics
5.5 Use in Patients with Reduced UGT1A1 Activity		
5.6 Embryo-Fetal Toxicity		13 NONCLINICAL TOXICOLOGY
6 ADVERSE REACTIONS		13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
6.1 Clinical Trials Experience		
6.2 Immunogenicity		14 CLINICAL STUDIES
7 DRUG INTERACTIONS		15 REFERENCES
7.1 UGT1A1 Inhibitors or Inducers		16 HOW SUPPLIED/STORAGE AND HANDLING
		17 PATIENT COUNSELING INFORMATION

\* Sections or subsections omitted from the full prescribing information are not listed.

# FULL PRESCRIBING INFORMATION

## WARNING: NEUTROPENIA AND DIARRHEA

- Severe neutropenia may occur. Withhold TRODELVY for absolute neutrophil count below 1500/mm<sup>3</sup> or neutropenic fever. Monitor blood cell counts periodically during treatment. Consider G-CSF for secondary prophylaxis. Initiate anti-infective treatment in patient with febrile neutropenia without delay [see Warnings and Precautions (5.1)].
- Severe diarrhea may occur. Monitor patients with diarrhea and give fluid and electrolytes as needed. Administer atropine, if not contraindicated, for early diarrhea of any severity. At the onset of late diarrhea, evaluate for infectious causes and, if negative, promptly initiate loperamide [see Warnings and Precautions (5.2)]. If severe diarrhea occurs, withhold TRODELVY until resolved to ≤ Grade 1 and reduce subsequent doses [see Dosage and Administration (2.3)].

## 1 INDICATIONS AND USAGE

TRODELVY is indicated for the treatment of adult patients with metastatic triple-negative breast cancer (mTNBC) who have received at least two prior therapies for metastatic disease.

This indication is approved under accelerated approval based on tumor response rate and duration of response [see Clinical Studies (14)]. Continued approval for this indication may be contingent upon verification and description of clinical benefit in confirmatory trials.

## 2 DOSAGE AND ADMINISTRATION

### 2.1 Important Use Information

**Do NOT substitute TRODELVY for or use with other drugs containing irinotecan or its active metabolite SN-38.**

### 2.2 Recommended Dose and Schedule

The recommended dose of TRODELVY is 10 mg/kg administered as an intravenous infusion once weekly on Days 1 and 8 of 21-day treatment cycles. Continue treatment until disease progression or unacceptable toxicity. Do not administer TRODELVY at doses greater than 10 mg/kg.

Administer TRODELVY as an intravenous infusion only. Do not administer as an intravenous push or bolus.

First infusion: Administer infusion over 3 hours. Observe patients during the infusion and for at least 30 minutes following the initial dose, for signs or symptoms of infusion-related reactions [see Warning and Precautions (5.3)].

Subsequent infusions: Administer infusion over 1 to 2 hours if prior infusions were tolerated. Observe patients during the infusion and for at least 30 minutes after infusion.

#### Premedication

Prior to each dose of TRODELVY, premedication for prevention of infusion reactions and prevention of chemotherapy-induced nausea and vomiting (CINV) is recommended.

- Premedicate with antipyretics, H1 and H2 blockers prior to infusion, and corticosteroids may be used for patients who had prior infusion reactions.
- Premedicate with a two or three drug combination regimen (e.g., dexamethasone with either a 5-HT<sub>3</sub> receptor antagonist or an NK<sub>1</sub> receptor antagonist, as well as other drugs as indicated).

### 2.3 Dose Modifications for Adverse Reactions

#### Infusion-related Reactions

Slow or interrupt the infusion rate of TRODELVY if the patient develops an infusion-related reaction. Permanently discontinue TRODELVY for life-threatening infusion-related reactions [see Warnings and Precautions (5.3)]

#### Dose Modifications for Adverse Reactions

Withhold or discontinue TRODELVY to manage adverse reactions as described in Table 1. Do not re-escalate the TRODELVY dose after a dose reduction for adverse reactions has been made.

**Table 1: Dose Modifications for Adverse Reactions**

Adverse Reaction	Occurrence	Dose Modification
<b>Severe Neutropenia</b> [see Warnings and Precautions (5.1)]		
Grade 4 neutropenia ≥7 days, OR Grade 3 febrile neutropenia (absolute neutrophil count <1000/mm <sup>3</sup> and fever ≥38.5°C), OR At time of scheduled treatment, Grade 3-4 neutropenia which delays dosing by 2 or 3 weeks for recovery to ≤ Grade 1	First	25% dose reduction and administer granulocyte-colony stimulating factor (G-CSF)
	Second	50% dose reduction
	Third	Discontinue treatment
At time of scheduled treatment, Grade 3-4 neutropenia which delays dosing beyond 3 weeks for recovery to ≤ Grade 1	First	Discontinue treatment
<b>Severe Non-Neutropenic Toxicity</b>		
Grade 4 non-hematologic toxicity of any duration, OR Any Grade 3-4 nausea, vomiting or diarrhea due to treatment that is not controlled with antiemetics and anti-diarrheal agents [see Warnings and Precautions (5.2, 5.4)], OR Other Grade 3-4 non-hematologic toxicity persisting >48 hours despite optimal medical management, OR At time of scheduled treatment, Grade 3-4 non-neutropenic hematologic or non-hematologic toxicity, which delays dose by 2 or 3 weeks for recovery to ≤ Grade 1	First	25% dose reduction
	Second	50% dose reduction
	Third	Discontinue treatment
In the event of Grade 3-4 non-neutropenic hematologic or non-hematologic toxicity, which does not recover to ≤ Grade 1 within 3 weeks	First	Discontinue treatment

# FULL PRESCRIBING INFORMATION

## 2.4 Preparation for Administration

### Reconstitution

- TRODELVY is a cytotoxic drug.
- Follow applicable special handling and disposal procedures<sup>1</sup>.
- Calculate the required dose (mg) of TRODELVY based on the patient's body weight at the beginning of each treatment cycle (or more frequently if the patient's body weight changed by more than 10% since the previous administration) [see *Dosage and Administration* (2.1)].
- Allow the required number of vials to warm to room temperature.
- Using a sterile syringe, slowly inject 20 mL of 0.9% Sodium Chloride Injection, USP, into each 180 mg TRODELVY vial. The resulting concentration will be 10 mg/mL.
- Gently swirl vials and allow to dissolve for up to 15 minutes. Do not shake. Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. The solution should be free of visible particulates, clear and yellow. Do not use the reconstituted solution if it is cloudy or discolored.
- Use immediately to prepare a diluted TRODELVY infusion solution.

### Dilution

- Calculate the required volume of the reconstituted TRODELVY solution needed to obtain the appropriate dose according to patient's body weight. Withdraw this amount from the vial(s) using a syringe. Discard any unused portion remaining in the vial(s).
- Slowly inject the required volume of reconstituted TRODELVY solution into a polypropylene (PP) infusion bag, to minimize foaming. Do not shake the contents.
- Adjust the volume in the infusion bag as needed with 0.9% Sodium Chloride Injection, USP, to obtain a concentration of 1.1 mg/mL to 3.4 mg/mL (total volume should not exceed 500 mL). For patients whose body weight exceeds 170 kg, divide the total dosage of TRODELVY equally between two 500 mL infusion bags and infuse sequentially via slow infusion.
- Only 0.9% Sodium Chloride Injection, USP, should be used since the stability of the reconstituted product has not been determined with other infusion-based solutions. Use the diluted solution in the infusion bag immediately. If not used immediately, the infusion bag containing TRODELVY solution can be stored refrigerated 2°C to 8°C (36°F to 46°F) for up to 4 hours. After refrigeration, administer diluted solution within 4 hours (including infusion time).

Do Not Freeze or Shake. Protect from Light.

### Administration

- Administer TRODELVY as an intravenous infusion. Protect infusion bag from light.
- An infusion pump may be used.
- Do not mix TRODELVY, or administer as an infusion, with other medicinal products.
- Upon completion of the infusion, flush the intravenous line with 20 mL 0.9% Sodium Chloride Injection, USP.

## 3 DOSAGE FORMS AND STRENGTHS

For injection: 180 mg off-white to yellowish lyophilized powder in a single-dose vial.

## 4 CONTRAINDICATIONS

TRODELVY is contraindicated in patients who have experienced a severe hypersensitivity reaction to TRODELVY [see *Warnings and Precautions* (5.3)].

## 5 WARNINGS AND PRECAUTIONS

### 5.1 Neutropenia

TRODELVY can cause severe or life-threatening neutropenia. Withhold TRODELVY for absolute neutrophil count below 1500/mm<sup>3</sup> on Day 1 of any cycle or neutrophil count below 1000/mm<sup>3</sup> on Day 8 of any cycle. Withhold TRODELVY for neutropenic fever. Dose modifications may be required due to neutropenia [see *Dosage and Administration* (2.3)].

Febrile neutropenia occurred in 6% (24/408) patients treated with TRODELVY, including 8% (9/108) patients with mTNBC after at least two prior therapies. Less than 1% (1/408) of patients had febrile neutropenia leading to permanent discontinuation.

The incidence of Grade 1-4 neutropenia was 64% in patients with mTNBC (n=108). In all patients treated with TRODELVY (n=408), the incidence of Grade 1-4 neutropenia was 54%; Grade 4 neutropenia occurred in 13%. Less than 1% (2/408) of patients permanently discontinued treatment due to neutropenia.

### 5.2 Diarrhea

TRODELVY can cause severe diarrhea. Withhold TRODELVY for Grade 3-4 diarrhea at the time of scheduled treatment administration and resume when resolved to ≤ Grade 1 [see *Dosage and Administration* (2.3)].

At the onset of diarrhea, evaluate for infectious causes and if negative, promptly initiate loperamide, 4 mg initially followed by 2 mg with every episode of diarrhea for a maximum of 16 mg daily. Discontinue loperamide 12 hours after diarrhea resolves. Additional supportive measures (e.g., fluid and electrolyte substitution) may also be employed as clinically indicated.

Patients who exhibit an excessive cholinergic response to treatment with TRODELVY (e.g., abdominal cramping, diarrhea, salivation, etc.) can receive appropriate premedication (e.g., atropine) for subsequent treatments.

Diarrhea occurred in 63% (68/108) of patients with mTNBC and 62% (254/408) of all patients treated with TRODELVY. In each population, events of Grade 3-4 occurred in 9% (10/108) of mTNBC patients and 9% (36/408) of all patients treated with TRODELVY. Four out of 408 patients (<1%) discontinued treatment because of diarrhea. Neutropenic colitis was observed in 2% (2/108) of patients in the mTNBC cohort and 1% of all patients treated with TRODELVY.

# FULL PRESCRIBING INFORMATION

## 5.3 Hypersensitivity

TRODELVY can cause severe and life-threatening hypersensitivity. Anaphylactic reactions have been observed in clinical trials with TRODELVY.

Hypersensitivity reactions within 24 hours of dosing occurred in 37% (151/408) of patient treated with TRODELVY. Grade 3-4 hypersensitivity occurred in 1% (6/408) of patients treated with TRODELVY. The incidence of hypersensitivity reactions leading to permanent discontinuation of TRODELVY was 1% (3/408).

Pre-infusion medication for patients receiving TRODELVY is recommended. Observe patients closely for infusion-related reactions during each TRODELVY infusion and for at least 30 minutes after completion of each infusion [see *Dosage and Administration* (2.3)]. Medication to treat such reactions, as well as emergency equipment, should be available for immediate use.

## 5.4 Nausea and Vomiting

TRODELVY is emetogenic. Nausea occurred in 69% (74/108) of patients with mTNBC and 69% (281/408) of all patients treated with TRODELVY. Grade 3 nausea occurred in 6% (7/108) and 5% (22/408) of these populations, respectively.

Vomiting occurred in 49% (53/108) of patients with mTNBC and 45% (183/408) of all patients treated with TRODELVY. Grade 3 vomiting occurred in 6% (7/108) and 4% (16/408) of these patients, respectively.

Premedicate with a two or three drug combination regimen (e.g., dexamethasone with either a 5-HT<sub>3</sub> receptor antagonist or an NK-1 receptor antagonist as well as other drugs as indicated) for prevention of chemotherapy-induced nausea and vomiting (CINV).

Withhold TRODELVY doses for Grade 3 nausea or Grade 3-4 vomiting at the time of scheduled treatment administration and resume with additional supportive measures when resolved to Grade ≤ 1 [see *Dosage and Administration* (2.3)].

Additional antiemetics and other supportive measures may also be employed as clinically indicated. All patients should be given take-home medications with clear instructions for prevention and treatment of nausea and vomiting.

## 5.5 Use in Patients with Reduced UGT1A1 Activity

Individuals who are homozygous for the uridine diphosphate-glucuronosyl transferase 1A1 (UGT1A1)\*28 allele are at increased risk for neutropenia and may be at increased risk for other adverse reactions following initiation of TRODELVY treatment.

In 84% (343/408) of patients who received TRODELVY (up to 10 mg/kg on Days 1 and 8 of a 21-day cycle) and had retrospective UGT1A1 genotype results available, the incidence of Grade 4 neutropenia was 26% (10/39) in patients homozygous for the UGT1A1\*28 allele, 13% (20/155) in patients heterozygous for the UGT1A1\*28 allele and 11% (16/149) in patients homozygous for the wild-type allele [see *Clinical Pharmacology* (12.5)].

Closely monitor patients with reduced UGT1A1 activity for severe neutropenia. The appropriate dose for patients who are homozygous for UGT1A1\*28 is not known and should be considered based on individual patient tolerance to treatment [see *Dosage and Administration* (2.3)].

## 5.6 Embryo-Fetal Toxicity

Based on its mechanism of action, TRODELVY can cause teratogenicity and/or embryo-fetal lethality when administered to a pregnant woman. TRODELVY contains a genotoxic component, SN-38, and targets rapidly dividing cells [see *Clinical Pharmacology* (12.1) and

*Nonclinical Toxicology* (13.1)]. Advise pregnant women and females of reproductive potential of the potential risk to a fetus. Advise females of reproductive potential to use effective contraception during treatment with TRODELVY and for 6 months after the last dose. Advise male patients with female partners of reproductive potential to use effective contraception during treatment with TRODELVY and for 3 months after the last dose [see *Use in Specific Populations* (8.1, 8.3)].

## 6 ADVERSE REACTIONS

The following adverse reactions are discussed in greater detail in other sections of the label:

- Neutropenia [see *Warnings and Precautions* (5.1)]
- Diarrhea [see *Warnings and Precautions* (5.2)]
- Hypersensitivity [see *Warnings and Precautions* (5.3)]
- Nausea and Vomiting [see *Warnings and Precautions* (5.4)]

### 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

The data described in the Warnings and Precautions section reflect exposure to TRODELVY as a single agent in a single-arm, open-label study (IMMU-132-01) in 408 patients with mTNBC and other malignancies who had received prior systemic therapeutic regimen for advanced disease. TRODELVY was administered as an intravenous infusion once weekly on Days 1 and 8 of 21-day treatment cycles at doses up to 10 mg/kg until disease progression or unacceptable toxicity.

The data in Table 2 reflect exposure to TRODELVY in a subset of 108 patients with mTNBC who had received at least two prior treatments for metastatic disease in study (IMMU-132-01). Patients received TRODELVY 10 mg/kg via intravenous infusion on Days 1 and 8 of 21-day treatment cycles until disease progression or unacceptable toxicity. The median treatment duration in these 108 patients was 5.1 months (range: 0-51 months).

Serious adverse reactions were reported in 31% of the patients. The most frequent serious adverse reactions (reported in >1% of the patients receiving TRODELVY) were febrile neutropenia (6%), vomiting (5%), nausea (3%), dyspnea (3%), diarrhea (4%), anemia (2%), pleural effusion, neutropenia, pneumonia, dehydration (each 2%).

TRODELVY was permanently discontinued for adverse reactions in 2% of patients. Adverse reactions leading to discontinuation were anaphylaxis, anorexia/fatigue, headache (each <1%, 1 patient for each event). Forty-five percent (45%) of patients experienced an adverse reaction leading to treatment interruption. The most common adverse reaction leading to treatment interruption was neutropenia (33%). Adverse reactions leading to dose reduction occurred in 33% of patients treated with TRODELVY, with 24% having one dose reduction and 9% with two dose reductions. The most common adverse reaction leading to dose reductions was neutropenia/febrile neutropenia.

Adverse reactions occurring in ≥10% of patients with mTNBC in the IMMU-132-01 study are summarized in Table 2.

# FULL PRESCRIBING INFORMATION

**Table 2: Adverse Reactions in ≥ 10% of Patients With mTNBC in IMMU-132-01**

Adverse Reaction	TRODELVY (n=108)	
	Grade 1-4 (%)	Grade 3-4 (%)
<b>Any adverse reaction</b>	100	71
<b>Gastrointestinal disorders</b>	95	21
Nausea	69	6
Diarrhea	63	9
Vomiting	49	6
Constipation	34	1
Abdominal pain <sup>i</sup>	26	1
Mucositis <sup>ii</sup>	14	1
<b>General disorders and administration site conditions</b>	77	9
Fatigue <sup>iii</sup>	57	8
Edema <sup>iv</sup>	19	0
Pyrexia	14	0
<b>Blood and lymphatic system disorders</b>	74	37
Neutropenia	64	43
Anemia	52	12
Thrombocytopenia	14	3
<b>Metabolism and nutrition disorders</b>	68	22
Decreased appetite	30	1
Hyperglycemia	24	4
Hypomagnesemia	21	1
Hypokalemia	19	2
Hypophosphatemia	16	9
Dehydration	13	5
<b>Skin and subcutaneous tissue disorders</b>	63	4
Alopecia	38	0
Rash <sup>v</sup>	31	3
Pruritis	17	0
Dry Skin	15	0
<b>Nervous system disorders</b>	56	4
Headache	23	1
Dizziness	22	0
Neuropathy <sup>vi</sup>	24	0
Dysgeusia	11	0
<b>Infections and infestations</b>	55	12
Urinary Tract Infection	21	3
Respiratory Infection <sup>vii</sup>	26	3
<b>Musculoskeletal and connective tissue disorders</b>	54	1
Back pain	23	0
Arthralgia	17	0
Pain in extremity	11	0
<b>Respiratory, thoracic and mediastinal disorders</b>	54	5
Cough <sup>viii</sup>	22	0
Dyspnea <sup>ix</sup>	21	3
<b>Psychiatric disorders</b>	26	1
Insomnia	13	0

Graded per NCI CTCAE v. 4.0

<sup>i</sup> Including abdominal pain, distention, pain (upper), discomfort, tenderness

<sup>ii</sup> Including stomatitis, esophagitis, and mucosal inflammation

<sup>iii</sup> Including fatigue and asthenia

<sup>iv</sup> Including edema; and peripheral, localized, and periorbital edema

<sup>v</sup> Including rash; maculopapular, erythematous, and generalized rash; dermatitis acneiform; skin disorder, irritation, and exfoliation

<sup>vi</sup> Including gait disturbance, hypoesthesia, muscular weakness, paresthesia, and peripheral and sensory neuropathy

<sup>vii</sup> Including lower and upper respiratory tract infection, pneumonia, influenza, viral upper respiratory infection, bronchitis, and respiratory syncytial virus infection

<sup>viii</sup> Includes cough and productive cough

<sup>ix</sup> Includes dyspnea and exertional dyspnea

**Table 3: Laboratory Abnormalities Observed in >10% of Patients While Receiving TRODELVY**

Laboratory Abnormality	TRODELVY (n=108)	
	All Grades (%)	Grade 3-4 (%)
<b>Hematology</b>		
Decreased hemoglobin	93	6
Decreased leukocytes	91	26
Decreased neutrophils	82	32
Increased activated partial thromboplastin time	60	12
Decreased platelets	30	3
<b>Chemistry</b>		
Increased alkaline phosphatase	57	2
Decreased magnesium	51	3
Decreased calcium	49	3
Increased glucose	48	3
Increased aspartate aminotransferase	45	3
Decreased albumin	39	1
Increased alanine aminotransferase	35	2
Decreased potassium	30	3
Decreased phosphate	29	5
Decreased sodium	25	4.7
Increased magnesium	24	4
Decreased glucose	19	2

## 6.2 Immunogenicity

As with all therapeutic proteins, there is potential for immunogenicity. The detection of antibody formation is highly dependent on the sensitivity and specificity of the assay. Additionally, the observed incidence of antibody (including neutralizing antibody) positivity in an assay may be influenced by several factors including assay methodology, sample handling, timing of sample collection, concomitant medications, and underlying disease. For these reasons, comparison of the incidence of antibodies in the studies described below with the incidence of antibodies in other studies or to other sacituzumab govitecan products may be misleading.

The analysis of immunogenicity of TRODELVY in serum samples from 106 patients with mTNBC was evaluated using an electrochemiluminescence (ECL)-based immunoassay to test for anti-sacituzumab govitecan-hzy antibodies. Detection of the anti-sacituzumab govitecan-hzy antibodies was done using a 3-tier approach: screen, confirm, and titer. Persistent anti-sacituzumab govitecan-hzy antibodies developed in 2% (2/106) of patients.

## 7 DRUG INTERACTIONS

### 7.1 Effect of Other Drugs on TRODELVY

#### UGT1A1 Inhibitors

Concomitant administration of TRODELVY with inhibitors of UGT1A1 may increase the incidence of adverse reactions due to potential increase in systemic exposure to SN-38 [see *Warning and Precaution (5.5) and Clinical Pharmacology (12.3, 12.5)*]. Avoid administering UGT1A1 inhibitors with TRODELVY.

#### UGT1A1 Inducers

Exposure to SN-38 may be substantially reduced in patients concomitantly receiving UGT1A1 enzyme inducers [see *Warning and Precaution (5.5) and Clinical Pharmacology (12.3, 12.5)*]. Avoid administering UGT1A1 inducers with TRODELVY.

# FULL PRESCRIBING INFORMATION

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

#### Risk Summary

Based on its mechanism of action, TRODELVY can cause teratogenicity and/or embryo-fetal lethality when administered to a pregnant woman. There are no available data in pregnant women to inform the drug-associated risk. TRODELVY contains a genotoxic component, SN-38, and is toxic to rapidly dividing cells [see *Clinical Pharmacology (12.1) and Nonclinical Toxicology (13.1)*]. Advise pregnant women and females of reproductive potential of the potential risk to a fetus.

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 – 4% and 15 – 20%, respectively.

#### Data

##### *Animal data*

There were no reproductive and developmental toxicology studies conducted with sacituzumab govitecan-hziy.

### 8.2 Lactation

#### Risk Summary

There is no information regarding the presence of sacituzumab govitecan-hziy or SN-38 in human milk, the effects on the breastfed child, or the effects on milk production. Because of the potential for serious adverse reactions in a breastfed child, advise women not to breastfeed during treatment and for 1 month after the last dose of TRODELVY.

### 8.3 Females and Males of Reproductive Potential

#### Pregnancy Testing

Verify the pregnancy status of females of reproductive potential prior to the initiation of TRODELVY.

#### Contraception

##### *Females*

TRODELVY can cause fetal harm when administered to a pregnant woman [see *Use in Specific Populations (8.1)*]. Advise females of reproductive potential to use effective contraception during treatment with TRODELVY and for 6 months after the last dose.

##### *Males*

Because of the potential for genotoxicity, advise male patients with female partners of reproductive potential to use effective contraception during treatment with TRODELVY and for 3 months after the last dose.

#### Infertility

##### *Females*

Based on findings in animals, TRODELVY may impair fertility in females of reproductive potential [see *Nonclinical Toxicology (13.1)*].

### 8.4 Pediatric Use

Safety and effectiveness of TRODELVY have not been established in pediatric patients.

### 8.5 Geriatric Use

Of the patients who received TRODELVY, 19/108 (18%) patients with mTNBC and 144/408 (35%) of all patients were  $\geq 65$  years old. No overall differences in safety and effectiveness were observed between these patients and younger patients.

### 8.6 Hepatic Impairment

No adjustment to the starting dose is required when administering TRODELVY to patients with mild hepatic impairment (bilirubin less than or equal to 1.5 ULN and AST/ALT  $< 3$  ULN).

The exposure of TRODELVY in patients with mild hepatic impairment (bilirubin less than or equal to ULN and AST greater than ULN, or bilirubin greater than 1.0 to 1.5 ULN and AST of any level; n=12) was similar to patients with normal hepatic function (bilirubin or AST less than ULN; n=45).

The safety of TRODELVY in patients with moderate or severe hepatic impairment has not been established. TRODELVY has not been tested in patients with serum bilirubin  $> 1.5$  ULN, or AST and ALT  $> 3$  ULN, or AST and ALT  $> 5$  ULN and associated with liver metastases.

No dedicated trial was performed to investigate the tolerability of TRODELVY in patients with moderate or severe hepatic impairment. No recommendations can be made for the starting dose in these patients.

## 10 OVERDOSAGE

In a clinical trial, planned doses of up to 18 mg/kg (approximately 1.8 times the maximum recommended dose of 10 mg/kg) of TRODELVY were administered. In these patients, a higher incidence of severe neutropenia was observed.


## 11 DESCRIPTION

Sacituzumab govitecan-hziy is a Trop-2 directed antibody and topoisomerase inhibitor conjugate, composed of the following three components:

- the humanized monoclonal antibody, hRS7 IgG1 $\kappa$  (also called sacituzumab), which binds to Trop-2 (the trophoblast cell-surface antigen-2);
- the drug SN-38, a topoisomerase inhibitor;
- a hydrolysable linker (called CL2A), which links the humanized monoclonal antibody to SN-38.

## FULL PRESCRIBING INFORMATION

The recombinant monoclonal antibody is produced by mammalian (murine myeloma) cells, while the small molecule components SN-38 and CL2A are produced by chemical synthesis. Sacituzumab govitecan-hziy contains on average 7 to 8 molecules of SN-38 per antibody molecule. Sacituzumab govitecan-hziy has a molecular weight of approximately 160 kilodaltons. Sacituzumab govitecan-hziy has the following chemical structure.



TRODELVY (sacituzumab govitecan-hziy) for injection is a sterile, preservative-free, off-white to yellowish lyophilized powder for intravenous use in a 50 mL clear glass single-dose vial, with a rubber stopper and crimp-sealed with an aluminum flip-off cap.

Each single-dose vial of TRODELVY delivers 180 mg sacituzumab govitecan-hziy, 77.3 mg 2-(N-morpholino) ethane sulfonic acid (MES), 1.8 mg polysorbate 80 and 154 mg trehalose dihydrate. Reconstitution with 20 mL of 0.9% Sodium Chloride Injection, USP, results in a concentration of 10 mg/mL with a pH of 6.5.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Sacituzumab govitecan-hziy is a Trop-2-directed antibody-drug conjugate. Sacituzumab is a humanized antibody that recognizes Trop-2. The small molecule, SN 38, is a topoisomerase I inhibitor, which is covalently attached to the antibody by a linker. Pharmacology data suggest that sacituzumab govitecan-hziy binds to Trop-2-expressing cancer cells and is internalized with the subsequent release of SN-38 via hydrolysis of the linker. SN-38 interacts with topoisomerase I and prevents re-ligation of topoisomerase I-induced single strand breaks. The resulting DNA damage leads to apoptosis and cell death. Sacituzumab govitecan-hziy decreased tumor growth in mouse xenograft models of triple-negative breast cancer.

### 12.2 Pharmacodynamics

Exposure-response relationships and the time course of pharmacodynamics response are unknown for sacituzumab govitecan-hziy.

### 12.3 Pharmacokinetics

The serum pharmacokinetics of sacituzumab govitecan-hziy and SN-38 were evaluated in a study in a population of mTNBC patients who received sacituzumab govitecan-hziy as a single agent at a dose of 10 mg/kg. The pharmacokinetic parameters of sacituzumab govitecan-hziy and free SN-38 are presented in Table 4.

**Table 4: Summary of Mean PK Parameters ( $\pm$ Standard Deviation) of Sacituzumab Govitecan-hziy and Free SN-38**

	Sacituzumab govitecan-hziy	Free SN-38
$C_{max}$ [ng/mL]	243,000 ( $\pm 45,600$ )	127 ( $\pm 60$ )
$AUC_{0-168}$ [h ng/mL]	5,210,000 ( $\pm 1,230,000$ )	3,900 ( $\pm 1,830$ )

$C_{max}$ : maximum plasma concentration  
 $AUC_{0-168}$ : area under plasma concentration curve through 168 hours

### Distribution

The mean volume of distribution for sacituzumab govitecan-hziy was 0.045 L/kg.

### Elimination

The mean half-life of sacituzumab govitecan-hziy and free SN-38 was 16 and 18 hours, respectively. The clearance of the sacituzumab govitecan-hziy was 0.002 L/h/kg.

### Metabolism

No metabolism studies with sacituzumab govitecan-hziy have been conducted. SN-38 (the small molecule moiety of sacituzumab govitecan-hziy) is metabolized via UGT1A1. The glucuronide metabolite of SN-38 (SN-38G) was detectable in the serum of patients.

### Specific Populations

Pharmacokinetic analyses in a limited number of patients with mTNBC ( $n = 57$ ) did not identify an effect of age or race on the pharmacokinetics of sacituzumab govitecan-hziy. Renal elimination is known to contribute minimally to the excretion of SN-38, the small molecule moiety of sacituzumab govitecan-hziy. There are no data on the pharmacokinetics of sacituzumab govitecan-hziy in patients with renal impairment or end-stage renal disease ( $CL_{cr} \leq 30$  mL/min).

The exposure of sacituzumab govitecan-hziy is similar in patients with mild hepatic impairment (bilirubin less than or equal to ULN and AST greater than ULN, or bilirubin greater than 1.0 to less than 1.5 ULN and AST of any level;  $n=12$ ) to patients with normal hepatic function (bilirubin or AST less than ULN;  $n=45$ ).

Sacituzumab govitecan-hziy exposure is unknown in patients with moderate or severe hepatic impairment. SN-38 exposure may be elevated in such patients due to decreased hepatic UGT1A1 activity.

### Drug Interaction Studies

No drug-drug interaction studies were conducted with sacituzumab govitecan-hziy or its components. Inhibitors or inducers of UGT1A1 are expected to increase or decrease SN-38 exposure, respectively [see *Drug Interactions* (7)].

## FULL PRESCRIBING INFORMATION

### 12.5 Pharmacogenomics

SN-38 is metabolized via UGT1A1 [see *Clinical Pharmacology* (12.3)]. Genetic variants of the UGT1A1 gene such as the UGT1A1\*28 allele lead to reduced UGT1A1 enzyme activity. Individuals who are homozygous for the UGT1A1\*28 allele are at increased risk for neutropenia from TRODELVY [see *Warnings and Precautions* (5.5)]. Approximately 20% of the Black or African American population, 10% of the White population, and 2% of the East Asian population are homozygous for the UGT1A1\*28 allele. Decreased function alleles other than UGT1A1\*28 may be present in certain populations.

## 13 NONCLINICAL TOXICOLOGY

### 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenicity studies have not been conducted with sacituzumab govitecan-hziy.

SN-38 was clastogenic in an *in vitro* mammalian cell micronucleus test in Chinese hamster ovary cells and was not mutagenic in an *in vitro* bacterial reverse mutation (Ames) assay.

Fertility studies with sacituzumab govitecan-hziy have not been conducted. In a repeat-dose toxicity study in cynomolgus monkeys, intravenous administration of sacituzumab govitecan-hziy on Day 1 and Day 4 resulted in endometrial atrophy, uterine hemorrhage, increased follicular atresia of the ovary, and atrophy of vaginal epithelial cells at doses  $\geq$  60 mg/kg ( $\geq$  6 times the human recommended dose of 10 mg/kg based on body weight).

## 14 CLINICAL STUDIES

The efficacy of TRODELVY was evaluated in study IMMU-132-01 (NCT01631552), a multicenter, single-arm, trial that enrolled 108 patients with metastatic triple-negative breast cancer (mTNBC) who had received at least two prior treatments for metastatic disease. Patients with bulky disease, defined as a mass  $>$ 7 cm, were not eligible. Patients with treated brain metastases not receiving high dose steroids ( $>$ 20 mg prednisone or equivalent) for at least four weeks were eligible. Patients with known Gilbert's disease were excluded.

Patients received TRODELVY 10 mg/kg intravenously on Days 1 and 8 of a 21-day treatment cycle. Patients were treated with TRODELVY until disease progression or intolerance to the therapy. Tumor imaging was obtained every 8 weeks, with confirmatory CT/MRI scans obtained 4-6 weeks after an initial partial or complete response, until progression requiring treatment discontinuation. Major efficacy outcome measures were investigator assessed overall response rate (ORR) using RECIST 1.1 and duration of response.

The median age was 55 years (range: 31 – 80 years); 87% of patients were younger than 65 years. The majority of patients were female (99%), and White (76%). At study entry, all patients had an ECOG performance status of 0 (29%) or 1 (71%). Seventy-six percent had visceral disease, 42% had hepatic metastases, 56% had lung/pleura metastases, and 2% had brain metastases. Twelve patients (11%) had Stage IV disease at the time of initial diagnosis.

The median number of prior systemic therapies received in the metastatic setting was 3 (range: 2 - 10). Prior chemotherapies in the metastatic setting included carboplatin or cisplatin (69%), gemcitabine (55%), paclitaxel or docetaxel (53%), capecitabine (51%), eribulin (45%), doxorubicin (24%), vinorelbine (16%), cyclophosphamide (19%), and ixabepilone (8%).

Overall, 98% of patients had received prior taxanes and 86% had received prior anthracyclines either in the (neo)adjuvant or metastatic setting.

Table 5 summarizes the efficacy results.

**Table 5: Efficacy Results for Patients With mTNBC in IMMU-132-01**

TRODELVY (N=108)	
<b>Overall Response Rate<sup>i</sup></b>	
ORR (95% CI)	33.3% (24.6, 43.1)
Complete response	2.8%
Partial response	30.6%
<b>Response duration<sup>i</sup></b>	
Number of responders	36
Median, Months (95% CI)	7.7 (4.9, 10.8)
Range, Months	1.9+, 30.4+
% with duration $\geq$ 6 months	55.6%
% with duration $\geq$ 12 months	16.7%

<sup>i</sup> investigator assessment  
CI: confidence interval  
+: denotes ongoing

## 15 REFERENCES

1. "OSHA Hazardous Drugs." OSHA. <http://www.osha.gov/SLTC/hazardousdrugs/index.html>.

## 16 HOW SUPPLIED/STORAGE AND HANDLING

TRODELVY (sacituzumab govitecan-hziy) for injection is a sterile, off-white to yellowish lyophilized powder in a single-dose vial. Each TRODELVY vial is individually boxed in a carton:

- NDC 55135-132-01 contains one 180 mg vial

Store vials in a refrigerator at 2°C to 8°C (36°F to 46°F) in the original carton to protect from light until time of reconstitution. Do not freeze.

TRODELVY is a cytotoxic drug. Follow applicable special handling and disposal procedures<sup>1</sup>.

# FULL PRESCRIBING INFORMATION

## 17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information)

### Neutropenia

Advise patients of the risk of neutropenia. Instruct patients to immediately contact their healthcare provider if they experience fever, chills, or other signs of infection [see Warnings and Precautions (5.1)].

### Diarrhea

Advise patients of the risk of diarrhea. Instruct patients to immediately contact their healthcare provider if they experience diarrhea for the first time during treatment; black or bloody stools; symptoms of dehydration such as lightheadedness, dizziness, or faintness; inability to take fluids by mouth due to nausea or vomiting; or inability to get diarrhea under control within 24 hours [see Warnings and Precautions (5.2)].

### Hypersensitivity

Inform patients of the risk of serious infusion reactions and anaphylaxis. Instruct patients to immediately contact their healthcare provider if they experience facial, lip, tongue, or throat swelling, urticaria, difficulty breathing, lightheadedness, dizziness, chills, rigors, wheezing, pruritus, flushing, rash, hypotension or fever, that occur during or within 24 hours following the infusion [see Warnings and Precautions (5.3)].

### Nausea/Vomiting

Advise patients of the risk of nausea and vomiting. Premedication according to established guidelines with a two or three drug regimen for prevention of chemotherapy-induced nausea and vomiting (CINV) is also recommended. Additional antiemetics, sedatives, and other supportive measures may also be employed as clinically indicated. All patients should receive take-home medications for preventing and treating delayed nausea and vomiting, with clear instructions. Instruct patients to immediately contact their healthcare provider if they experience uncontrolled nausea or vomiting [see Warnings and Precautions (5.4)].

### Embryo-Fetal Toxicity

Advise female patients to contact their healthcare provider if they are pregnant or become pregnant. Inform female patients of the risk to a fetus and potential loss of the pregnancy [see Use in Specific Populations (8.1)].

### Contraception

Advise female patients of reproductive potential to use effective contraception during treatment and for 6 months after the last dose of TRODELVY [see Use in Specific Populations (8.3)].

Advise male patients with female partners of reproductive potential to use effective contraception during treatment and for 3 months after the last dose of TRODELVY [see Use in Specific Populations (8.3)].

### Lactation

Advise women not to breastfeed during treatment and for 1 month after the last dose of TRODELVY [see Use in Specific Populations (8.2)].

### Infertility

Advise females of reproductive potential that TRODELVY may impair fertility [see use in Specific Populations (8.3)].

Manufactured by: Immunomedics, Inc., 300 The American Road, Morris Plains, NJ 07950, USA  
U.S. License No. 1737

# PATIENT INFORMATION

## TRODELVY™ (troh-DELL-vee) (sacituzumab govitecan-hziy) injection

### What is the most important information I should know about TRODELVY?

#### TRODELVY can cause serious side effects, including:

- **Low white blood cell count (neutropenia).** Low white blood cell counts are common with TRODELVY and can sometimes be severe and lead to infections that can be life-threatening. Your healthcare provider should check your blood cell counts during treatment with TRODELVY. If your white blood cell count is too low, your healthcare provider may need to lower your dose of TRODELVY, give you a medicine to help prevent low blood cell count with future doses of TRODELVY, or in some cases may stop TRODELVY. Your healthcare provider may need to give you antibiotic medicines if you develop fever while your white blood cell count is low. **Call your healthcare provider right away if you develop any of the following signs of infection during treatment with TRODELVY:**

- fever
- chills
- cough
- shortness of breath
- burning or pain when you urinate

- **Severe diarrhea.** Diarrhea is common with TRODELVY and can also be severe. Your healthcare provider should monitor you for diarrhea and give you medicine as needed to help control your diarrhea. If you lose too much body fluids (dehydration) your healthcare provider may need to give you fluids and electrolytes to replace body salts. If diarrhea happens later in your treatment, your healthcare provider may check you to see if the diarrhea may be caused by an infection. Your healthcare provider may decrease your dose or stop TRODELVY if your diarrhea is severe and cannot be controlled with anti-diarrheal medicines.

#### Call your healthcare provider right away:

- the first time that you get diarrhea during treatment with TRODELVY
- if you have black or bloody stools
- if you have symptoms of losing too much body fluid (dehydration) and body salts, such as lightheadedness, dizziness or faintness
- if you are unable to take fluids by mouth due to nausea or vomiting
- if you are not able to get your diarrhea under control within 24 hours

### What is TRODELVY?

TRODELVY is a prescription medicine used to treat adults with breast cancer that is:

- estrogen and progesterone hormone receptor (HR) negative, and human epidermal growth factor receptor 2 (HER2)-negative (also called triple-negative breast cancer), **and**
- that has spread to other parts of the body metastatic, **and**
- who previously received at least two therapies for metastatic disease

It is not known if TRODELVY is safe and effective in people with moderate or severe liver problems.

It is not known if TRODELVY is safe and effective in children.

**Do not receive TRODELVY if you have had a severe allergic reaction to TRODELVY.** Ask your healthcare provider if you are not sure.

## PATIENT INFORMATION

### Before receiving TRODELVY, tell your healthcare provider about all of your medical conditions, including if you:

- have been told that you carry a gene for uridine diphosphate-glucuronosyl transferase A1 (UGT1A1)\*28. People who carry this gene have an increased risk of getting side effects with TRODELVY, especially low white blood cell counts. **See “What is the most important information I should know about TRODELVY?”**
- have liver problems.
- are pregnant or plan to become pregnant. TRODELVY can harm your unborn baby. Your healthcare provider should check to see if you are pregnant before you start receiving TRODELVY.
  - Females who can become pregnant should use effective birth control during treatment and for 6 months after your last dose of TRODELVY. Talk to your healthcare provider about birth control choices that may be right for you during this time.
  - Males with a female partner who can become pregnant should use effective birth control during treatment and for 3 months after your last dose of TRODELVY.
  - Tell your healthcare provider right away if you or your partner become pregnant during treatment with TRODELVY.
- are breastfeeding or plan to breastfeed. It is not known if TRODELVY passes into your breastmilk and can harm your baby. Do not breastfeed during treatment and for 1 month after your last dose of TRODELVY.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Certain medicines may affect the way TRODELVY works.

### How will I receive TRODELVY?

- Your healthcare provider will give you TRODELVY into your vein through an intravenous (IV) line.
- TRODELVY is given 1 time each week, on Day 1 and on Day 8 of a 21-day treatment cycle.
- You will receive the first dose of TRODELVY over 3 hours. If you tolerate the first dose well, future doses may be given over 1 to 2 hours.
- Before each dose of TRODELVY, you will receive medicines to help prevent infusion reactions, and nausea and vomiting.
- You will be monitored for side effects during and for at least 30 minutes after you receive each infusion of TRODELVY.
- Your healthcare provider may slow down or temporarily stop your infusion of TRODELVY if you have an infusion-related reaction, or permanently stop TRODELVY if you have a life-threatening infusion-related reaction.
- Your healthcare provider will decide how long you will continue to receive TRODELVY.

### What are the possible side effects of TRODELVY?

TRODELVY can cause serious side effects, including:

#### • See “What is the most important information I should know about TRODELVY?”

- **Severe and life-threatening allergic reactions.** TRODELVY can cause severe and life-threatening allergic reactions during infusion (infusion-related reactions). Tell your healthcare provider or nurse right away if you get any of the following symptoms of an allergic reaction during an infusion of TRODELVY or within 24 hours after you receive a dose of TRODELVY:

- swelling of your face, lips, tongue, or throat
- difficulty breathing or wheezing
- hives
- lightheadedness, dizziness, feeling faint or pass out
- skin rash or flushing of your skin
- chills or shaking chills (rigors)
- fever

- **Nausea and vomiting.** Nausea and vomiting are common with TRODELVY and can sometimes be severe. Before each dose of TRODELVY, you will receive medicines to help prevent nausea and vomiting. You should be given medicines to take home with you, along with instructions about how to take them to help prevent and treat any nausea and vomiting after you receive TRODELVY. Call your healthcare provider right away if you have nausea or vomiting that is not controlled with the medicines prescribed for you. Your healthcare provider may decide to decrease your dose or stop TRODELVY if your nausea and vomiting is severe and cannot be controlled with anti-nausea medicines.

The most common side effects of TRODELVY include:

- tiredness
- decreased red blood cell count
- hair loss
- constipation
- rash. **See “Severe and life-threatening allergic reactions” above.**
- decreased appetite
- stomach-area (abdomen) pain

TRODELVY may cause fertility problems in females, which could affect your ability to have a baby. Talk to your healthcare provider if fertility is a concern for you.

These are not all of the possible side effects of TRODELVY.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

### General information about the safe and effective use of TRODELVY.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. You can ask your pharmacist or healthcare provider for information about TRODELVY that is written for health professionals.

### What are the ingredients in TRODELVY?

**Active ingredient:** sacituzumab govitecan-hziy

**Inactive ingredients:** 2-(N-morpholino) ethane sulfonic acid (MES), polysorbate 80 and trehalose dihydrate

Manufactured by: Immunomedics, Inc., 300 The American Road, Morris Plains, NJ 07950, USA

U.S. License No. 1737

For more information about TRODELVY, go to [www.TRODELVY.com](http://www.TRODELVY.com) or call 1-888-983-4668.

The Patient Information has been approved by the U.S. Food and Drug Administration.

Issued: April 2020





**Immunomedics**

©2020 Immunomedics, Inc  
All Rights Reserved. Apr 2020  
2020-US-TROT-00005