

## ACTIVASE STROKE ASSESSMENT EDUCATION GUIDE

### Indication

Activase® (alteplase) is indicated for the treatment of acute ischemic stroke (AIS). Exclude intracranial hemorrhage as the primary cause of stroke signs and symptoms prior to initiation of treatment. Initiate treatment as soon as possible but within 3 hours after symptom onset.

### Important Safety Information

#### Contraindications

Do not administer Activase to treat acute ischemic stroke in the following situations in which the risk of bleeding is greater than the potential benefit: current intracranial hemorrhage (ICH); subarachnoid hemorrhage; active internal bleeding; recent (within 3 months) intracranial or intraspinal surgery or serious head trauma; presence of intracranial conditions that may increase the risk of bleeding (e.g., some neoplasms, arteriovenous malformations, or aneurysms); bleeding diathesis; and current severe uncontrolled hypertension.

**Please see select Important Safety Information throughout and the accompanying full Prescribing Information.**



# Assessing and treating AIS

A resource for managing AIS, from presentation through Activase administration

STROKE ASSESSMENT

[Find details about:](#)

TREATMENT

**ACTIVASE<sup>®</sup>**  
**ALTEPLASE**  
A RECOMBINANT TISSUE PLASMINOGEN ACTIVATOR

Please see select Important Safety Information on pages 19-21 and the accompanying full Prescribing Information.

# Clinical presentation of acute ischemic stroke (AIS)

## Common symptoms of anterior circulation stroke include<sup>1\*</sup>:

- Aphasia
- Disturbed consciousness
- Dysarthria
- Facial palsy
- Hemisensory deficits
- Unilateral motor deficit

## TREATMENT

## Some symptoms of posterior circulation stroke include<sup>2,3\*</sup>:

- Diplopia
- Dizziness
- Dysarthria
- Dysphagia
- Dystaxia or ataxia

## Conditions that may mimic stroke<sup>4</sup>

- Bell's Palsy
- Brain abscess or infection
- Complicated migraine
- Conversion disorder/psychogenic conditions
- Hypertensive encephalopathy
- Hypoglycemia
- Seizures
- Tumor

\*The symptoms listed for anterior and posterior strokes are not exhaustive, nor are they mutually exclusive. Symptoms alone are not sufficient to diagnose the location of the occlusion, which should be confirmed using brain imaging.<sup>1</sup>

# National Institutes of Health Stroke Scale (NIHSS)<sup>5</sup>

## STROKE ASSESSMENT

3

## NIH STROKE SCALE

## TREATMENT

CATEGORY	DESCRIPTION	SCORE
<b>1a. Level of Consciousness (LOC)</b> (Is the patient alert, drowsy, etc)	Alert; keenly responsive Arousable by minor stimulation Arousable by repeated or strong stimulation Unresponsive or reflex responses only	0 1 2 3
<b>1b. LOC Questions</b> (Ask the patient the month and his or her age. Answer must be correct)	Answers both correctly Answers one correctly Both incorrect	0 1 2
<b>1c. LOC Commands</b> (Ask patient to open/close eyes and then grip/release nonparetic hand)	Performs both correctly Performs one correctly Both incorrect	0 1 2
<b>2. Best Gaze</b> (Only horizontal movement tested. Oculo-cephalic reflex is OK, but not calorics. Eyes open—patient follows finger or face.)	Normal Partial gaze palsy Forced deviation	0 1 2
<b>3. Visual Fields</b> (Test by confrontation. Introduce visual stimulus to patient's upper and lower field quadrants.)	No visual loss Partial hemianopia Complete hemianopia Bilateral hemianopia	0 1 2 3
<b>4. Facial Palsy</b> (Ask patient to show teeth/smile, raise eyebrows, and close eyes.)	Normal Minor paralysis Partial paralysis Complete paralysis	0 1 2 3
<b>5. Motor Arm</b> (Alternately position the patient's legs. Extend each leg 30 degrees, always while supine).	No drift Drift Some effort against gravity No effort against gravity No movement Amputation or joint fusion	0 1 2 3 4 UN



# NIHSS<sup>5</sup> (cont'd)

## STROKE ASSESSMENT

3

### NIH STROKE SCALE

## TREATMENT

CATEGORY	DESCRIPTION	SCORE
<b>6. Motor Leg</b> (Alternately position the patient's legs. Extend each leg 30 degrees, always while supine).	No drift Drift Some effort against gravity No effort against gravity No movement Amputation or joint fusion	0 1 2 3 4 UN
<b>7. Limb Ataxia</b> (Finger-nose, heel-shin tests done on both sides.)	Absent Present in one limb Present in two limbs Amputation or joint fusion	0 1 2 UN
<b>8. Sensory</b> (Use a pinprick to face, arms, trunk, and legs—compare side to side. Assess patient's awareness of being touched.)	Normal Mild-to-moderate loss Severe or total loss	0 1 2
<b>9. Best Language</b> (Ask patient to name items, describe a picture, read a sentence; intubated patients should write responses.)	No aphasia Mild-to-moderate aphasia Severe aphasia Mute, global aphasia	0 1 2 3
<b>10. Dysarthria</b> (Evaluate speech clarity by asking patient to repeat listed words.)	Normal articulation Mild-to-moderate dysarthria Severe dysarthria Intubated or other physical barrier	0 1 2 UN
<b>11. Extinction and Inattention</b> (Use information from prior testing to identify neglect.)	No abnormality Partial inattention or extinction Profound hemi-inattention or extinction	0 1 2



# Patients with AIS may initially be misdiagnosed<sup>6</sup>

STROKE ASSESSMENT

4

INDIVIDUALIZED  
DISABILITY ASSESSMENT

TREATMENT

A retrospective chart review found 1 in 5 ischemic strokes was initially misdiagnosed (n=103/465)

## Not all disability deficits may be detected by the NIHSS

- Although the National Institutes of Health Stroke Scale (NIHSS) is a useful tool for rating the level of neurological deficit due to stroke, it does not measure all deficits, nor does it fully assess the impact of disability on a patient's daily living or activity.<sup>7-9</sup> Use of the NIHSS should not substitute for a comprehensive neurological examination<sup>7-9</sup>
- The NIHSS is recommended by the 2019 American Heart Association/American Stroke Association (AHA/ASA) to quantify the degree of neurological deficit with accuracy and reliability<sup>7</sup>
- The NIHSS was originally developed for stroke research trials<sup>10</sup>
- Studies indicate that the NIHSS grades lesion-specific deficits unevenly<sup>9</sup>



# Consider how individual deficits are potentially disabling:

STROKE ASSESSMENT

4

INDIVIDUALIZED  
DISABILITY ASSESSMENT

TREATMENT

While the National Institutes of Health Stroke Scale (NIHSS) can rate the level of neurological deficit due to stroke, it may not detect all deficits. For each patient, all neurological deficits present at the time of the treatment decision should be considered in the context of individual risk and benefit, as well as the patient's baseline functional status.<sup>7</sup>

Activase® (alteplase) clinical trials enrolled patients with a measurable neurological deficit, defined as impairment of language, motor function, cognition, gaze, vision, or neglect.<sup>8,11</sup>

## Deficits considered by the AHA/ASA to be disabling<sup>8</sup>:

- Complete hemianopsia ( $\geq 2$  on NIHSS question 3)
- Severe aphasia ( $\geq 2$  on NIHSS question 9)
- Visual or sensory extinction ( $\geq 1$  on NIHSS question 11)
- Any weakness limiting sustained effort against gravity ( $\geq 2$  on NIHSS question 6 or 7)
- Any deficits that lead to a total NIHSS score  $> 5$
- Any remaining deficit the patient or practitioner considers potentially disabling (clinical judgment is required)



# Determine if your patients' deficits are potentially disabling

STROKE ASSESSMENT

4

INDIVIDUALIZED  
DISABILITY ASSESSMENT

TREATMENT

## Consider the following:

- Is your patient able to ambulate independently?
- Are they able to speak and understand?

## Consider asking each patient the following questions:

- Will you be able to return to work as normal?
- Will you be able to perform your everyday activities and hobbies?
- Do you consider any of your symptoms to be disabling?
- What is your dominant hand? Will weakness of your hand affect your daily life?
- Do you think your quality of life will be impacted?





# Quantify the level of disability

## STROKE ASSESSMENT

5

### MODIFIED RANKIN SCALE

#### TREATMENT

Measurement of a patient's ability to perform activities of daily living, including instrumental activities, should be a treatment consideration.<sup>12</sup> The modified Rankin Scale measures the ability of a patient to function independently without assistance.<sup>13</sup>

### Modified Rankin Scale<sup>13</sup>:

SCORE	DESCRIPTION
0	No symptoms at all
1	No significant disability despite symptoms; able to carry out all usual duties and activities
2	Slight disability; unable to carry out all previous activities but able to look after own affairs without assistance
3	Moderate disability; requiring some help but able to walk without assistance
4	Moderately severe disability; unable to walk without assistance and unable to attend to own bodily needs without assistance
5	Severe disability; bedridden, incontinent, and requiring constant nursing care and attention
6	Dead

# AHA/ASA Guideline for use of alteplase

## STROKE ASSESSMENT

### TREATMENT

6

#### AHA/ASA GUIDELINE FOR USE OF ALTEPLASE

### The AHA/ASA 2019 update to the 2018 Guidelines recommends IV alteplase use for management of AIS<sup>7</sup>

IV alteplase (0.9 mg/kg, maximum dose 90 mg over 60 minutes with initial 10% of dose given as bolus over 1 minute) is recommended for selected patients who may be treated within 3 hours of ischemic stroke symptom onset or patient last known well or at baseline state (**Class I; Level of Evidence A**).

In patients eligible for IV alteplase, benefit of therapy is time dependent, and treatment should be initiated as quickly as possible (**Class I; Level of Evidence A**).

### Activase is also recommended prior to mechanical thrombectomy

AHA/ASA 2019 update to the 2018 Guidelines<sup>5</sup>

- Patients eligible for IV alteplase should receive IV alteplase even if mechanical thrombectomy is being considered (**Class 1; Level of Evidence A**).

SVIN 2015 Recommendations<sup>14</sup>

- Endovascular mechanical thrombectomy, in addition to treatment with IV rtPA in eligible patients, is recommended for anterior circulation LVO ischemic strokes in patients presenting within 6 hours after symptom onset (**Class 1; Level of Evidence A**).

**ACTIVASE<sup>®</sup>**  
**ALTEPLASE**  
A RECOMBINANT TISSUE PLASMINOGEN ACTIVATOR

LVO=large vessel occlusion  
rtPA=recombinant tissue plasminogen activator  
SVIN-Society of Vascular and Interventional Neurology

Please see select Important Safety Information on pages 19-21 and the accompanying full Prescribing Information.

# Patient selection for Activase therapy<sup>11</sup>

## STROKE ASSESSMENT

## TREATMENT

7

## APPROPRIATE PATIENT SELECTION

- Activase is indicated for the treatment of AIS
- Exclude intracranial hemorrhage as the primary cause of stroke signs and symptoms prior to initiation of treatment
- Initiate treatment as soon as possible but within 3 hours after symptom onset

## Contraindications to Activase® (alteplase) therapy<sup>11</sup>

Do not administer Activase to treat AIS in the following situations in which the risk of bleeding is greater than the potential benefit:

- ☐ Current intracranial hemorrhage
- ☐ Subarachnoid hemorrhage
- ☐ Active internal bleeding
- ☐ Recent (within 3 months) intracranial or intraspinal surgery or serious head trauma
- ☐ Presence of intracranial conditions that may increase the risk of bleeding (eg, some neoplasms, arteriovenous malformations, or aneurysms)
- ☐ Bleeding diathesis\*
- ☐ Current severe uncontrolled hypertension

\*The 2019 update to the 2018 AHA/ASA Guideline advises against treatment with IV alteplase in patients<sup>7</sup>:

- With a platelet count <100,000/mm<sup>3</sup>, INR >1.7, aPTT >40 seconds, or PT >15 seconds
- Who have a history of warfarin use and an INR >1.7 and/or a PT >15 seconds
- Who have received a treatment dose of low-molecular-weight heparin within the previous 24 hours
- Who are taking direct thrombin inhibitors or direct factor Xa inhibitors, unless the laboratory tests are normal or the patient has not received a dose of these agents for >48 hours

Please see select Important Safety Information on pages 19-21 and the accompanying full Prescribing Information.

**ACTIVASE®**  
**ALTEPLASE**  
A RECOMBINANT TISSUE PLASMINOGEN ACTIVATOR

# NIH-recommended door-to-needle (DTN) times for AIS

STROKE ASSESSMENT

Since 2015, the Joint Commission has required DTN of  $\leq 60$  minutes in 50% of all eligible AIS patients receiving Activase® (alteplase)<sup>15</sup>


Target: Stroke has established a more aggressive goal<sup>16</sup>:

- DTN within 60 minutes in at least 75% of patients
- DTN within 45 minutes in at least 50% of patients

TREATMENT

7 APPROPRIATE PATIENT SELECTION

## DTN $\leq 60$ minutes<sup>16</sup>



## DTN $\leq 45$ minutes<sup>16</sup>



\*Initiate treatment with Activase as soon as possible but within 3 hours after symptom onset.<sup>11</sup>

ED=emergency department; NIH=National Institutes of Health.

Please see select Important Safety Information on pages 19-21 and the accompanying full Prescribing Information.

**ACTIVASE®**  
**ALTEPLASE**  
A RECOMBINANT TISSUE PLASMINOGEN ACTIVATOR

# FDA-approved dosing of Activase for AIS<sup>11</sup>

STROKE ASSESSMENT

TREATMENT

8

DOSING

The recommended dose of Activase is 0.9 mg/kg (not to exceed 90-mg total dose) infused intravenously over 60 minutes with 10% of the total dose administered as an initial bolus over 1 minute



**ACTIVASE®**  
**ALTEPLASE**  
A RECOMBINANT TISSUE PLASMINOGEN ACTIVATOR

FDA=Food and Drug Administration

Please see select Important Safety Information on pages 19-21 and the accompanying full Prescribing Information.

# Blood glucose management in patients with AIS<sup>7</sup>

STROKE ASSESSMENT

TREATMENT

## Recommendations from AHA/ASA 2019 update to the 2018 Guidelines

- Evidence indicates that persistent in-hospital hyperglycemia during the first 24 hours after AIS is associated with worse outcomes than normoglycemia, and thus, it is reasonable to treat hyperglycemia to achieve blood glucose levels in a range of 140 to 180 mg/dL and to closely monitor to prevent hypoglycemia (**Class IIa; Level of Evidence C-LD**)
- Hypoglycemia (blood glucose <60 mg/dL) should be treated in patients with AIS (**Class I; Level of Evidence C-LD**)

9

BLOOD GLUCOSE  
MANAGEMENT

**ACTIVASE<sup>®</sup>**  
**ALTEPLASE**  
A RECOMBINANT TISSUE PLASMINOGEN ACTIVATOR

Class IIa = procedure can be considered effective (moderate strength; benefit can be greater than risk);

Level C-LD = limited data available to support the recommendation.

Please see select Important Safety Information on pages 19-21 and the accompanying full Prescribing Information.

# Blood pressure (BP) management in patients with AIS<sup>7</sup>

STROKE ASSESSMENT

TREATMENT

## Recommendations from AHA/ASA 2019 update to the 2018 Guidelines

If patients are otherwise eligible for acute reperfusion except that BP is  $>185/110$  mm Hg, administer:

- Labetalol 10-20 mg IV over 1-2 minutes, may repeat 1 time; or
- Nicardipine 5 mg/h IV, titrate up by 2.5 mg/h every 5-15 minutes, maximum 15 mg/h, when desired BP reached, adjust to maintain proper BP limits; or
- Clevidipine 1-2 mg/h, titrate by doubling the dose every 2-5 min until desired BP reached, maximum 21 mg/h; or
- Other agents (eg, hydralazine, enalaprilat) may also be considered

If BP is not maintained  $\leq 185/110$  mm Hg, do not administer alteplase.

## BP management in patients with AIS<sup>7</sup>

- In patients with AIS, early treatment of hypertension is indicated when required by comorbid conditions (eg, concomitant acute coronary event, acute heart failure, aortic dissection, postthrombolysis symptomatic intracranial hemorrhage, or preeclampsia/eclampsia). Lowering BP initially by 15% is probably safe (**Class I; Level of Evidence C-EO**)
- In patients with BP  $\geq 220/120$  mm Hg who did not receive IV alteplase or endovascular therapy and have no comorbid conditions requiring acute antihypertensive treatment, the benefit of initiating or reinitiating treatment of hypertension within the first 48 to 72 hours is uncertain. It might be reasonable to lower BP by 15% during the first 24 hours after onset of stroke (**Class IIb; Level of Evidence C-EO**)

Level C-EO=consensus of expert opinion based on clinical experience.

Please see select Important Safety Information on pages 19-21 and the accompanying full Prescribing Information.

**ACTIVASE<sup>®</sup>**  
**ALTEPLASE**  
A RECOMBINANT TISSUE PLASMINOGEN ACTIVATOR



# Overall management of BP in patients treated with Activase<sup>7</sup>

STROKE ASSESSMENT

TREATMENT

## Recommendations from AHA/ASA 2019 update to the 2018 Guidelines

To maintain BP  $\leq$ 180/105 mm Hg, monitor BP:

- Every 15 minutes for 2 hours from the start of alteplase therapy;
- Then every 30 minutes for 6 hours; and
- Then every hour for 16 hours

If systolic BP is >180-230 mm Hg or diastolic BP is >105-120 mm Hg, administer:

- Labetalol 10 mg IV followed by continuous IV infusion 2-8 mg/min; or
- Nicardipine 5 mg/h IV, titrate up to desired effect by 2.5 mg/h every 5-15 minutes (maximum 15 mg/h); or
- Clevidipine 1-2 mg/h IV, titrate by doubling the dose every 2-5 min until desired BP reached (maximum 21 mg/h)

If BP is not controlled or diastolic BP >140 mm Hg, consider IV sodium nitroprusside.

10

BLOOD PRESSURE  
MANAGEMENT

**ACTIVASE<sup>®</sup>**  
**ALTEPLASE**  
A RECOMBINANT TISSUE PLASMINOGEN ACTIVATOR

Please see select Important Safety Information on pages 19-21 and the accompanying full Prescribing Information.





# Patient monitoring during Activase administration

STROKE ASSESSMENT

TREATMENT

## Monitor BP every 15 minutes during the 1-hour infusion<sup>7,11</sup>

- Once IV alteplase is given, the BP must be maintained at or below 180/105 mm Hg to limit the risk of intracranial hemorrhage
- Administer antihypertensive medications to maintain BP at or below these levels

## Perform neurological assessments<sup>5</sup>

The use of a stroke rating scale, preferably the NIHSS, is recommended

- Repeat every 15 minutes during the 1-hour infusion to monitor for neurological deterioration

## Check for major and/or minor bleeding<sup>11,17</sup>

All body secretions should be tested for occult blood

- Major bleeding: intracranial, retroperitoneal, gastrointestinal, or genitourinary hemorrhages
- Minor bleeding: gums, venipuncture sites, hematuria, hemoptysis, skin hematomas, or ecchymosis
- Arterial and venous punctures should be minimized and checked frequently deterioration

Discontinue infusion and obtain an emergency CT scan if the patient develops severe headache, acute hypertension, nausea, or vomiting or has a worsening neurological examination<sup>5</sup>

## Monitor for signs of hypersensitivity<sup>11</sup>

If signs of hypersensitivity occur, such as anaphylactoid reaction or development of angioedema, discontinue the Activase infusion and promptly institute appropriate therapy

Please see select Important Safety Information on pages 19-21 and the accompanying full Prescribing Information.

11

PATIENT MONITORING

**ACTIVASE<sup>®</sup>**  
**ALTEPLASE**  
A RECOMBINANT TISSUE PLASMINOGEN ACTIVATOR



# Patient monitoring post-Activase administration

STROKE ASSESSMENT

TREATMENT

## Continue to monitor for neurological deterioration<sup>7</sup>

- Every 15 minutes for the first hour after the infusion is stopped
- Every 30 minutes for the next 6 hours
- Hourly from the eighth postinfusion hour until 24 hours after the infusion is stopped

## Continue to check for major and/or minor bleeding<sup>17</sup>

## Continue to monitor and control BP<sup>7</sup>

- Every 15 minutes for the first hour after the infusion is stopped
- Every 30 minutes for the next 6 hours
- Hourly from the eighth postinfusion hour until 24 hours after the infusion is stopped

## Obtain a follow-up CT scan or MRI<sup>7</sup>

At 24 hours before starting anticoagulants or antiplatelet agents.

## Continue to monitor for hypersensitivity<sup>11</sup>

Please see select Important Safety Information on pages 19-21 and the accompanying full Prescribing Information.

11

PATIENT MONITORING

**ACTIVASE<sup>®</sup>**  
**ALTEPLASE**  
A RECOMBINANT TISSUE PLASMINOGEN ACTIVATOR



# Indication and Important Safety Information

## STROKE ASSESSMENT

### Indication

Activase® (alteplase) is indicated for the treatment of acute ischemic stroke (AIS). Exclude intracranial hemorrhage as the primary cause of stroke signs and symptoms prior to initiation of treatment. Initiate treatment as soon as possible but within 3 hours after symptom onset.

## TREATMENT

### Important Safety Information

#### Contraindications

Do not administer Activase to treat acute ischemic stroke in the following situations in which the risk of bleeding is greater than the potential benefit: current intracranial hemorrhage (ICH); subarachnoid hemorrhage; active internal bleeding; recent (within 3 months) intracranial or intraspinal surgery or serious head trauma; presence of intracranial conditions that may increase the risk of bleeding (e.g., some neoplasms, arteriovenous malformations, or aneurysms); bleeding diathesis; and current severe uncontrolled hypertension.

#### Warnings and Precautions

##### *Bleeding*

Activase can cause significant, sometimes fatal internal or external bleeding, especially at arterial and venous puncture sites. Avoid intramuscular injections and trauma to the patient. Perform venipunctures carefully and only as required. Fatal cases of hemorrhage associated with traumatic intubation in patients administered Activase have been reported. The concomitant administration of heparin and aspirin with and following infusions of Activase for the treatment of AIS during the first 24 hours after symptom onset has not been investigated.

Because heparin, aspirin, or Activase may cause bleeding complications, carefully monitor for bleeding, especially at arterial puncture sites. Hemorrhage can occur 1 or more days after administration of Activase, while patients are still receiving anticoagulant therapy. If serious bleeding occurs, terminate the Activase infusion, and treat properly.

12

## IMPORTANT SAFETY INFORMATION

**ACTIVASE®**  
**ALTEPLASE**  
A RECOMBINANT TISSUE PLASMINOGEN ACTIVATOR



# Important Safety Information (*cont'd*)

STROKE ASSESSMENT

TREATMENT

## Warnings and Precautions

### ***Bleeding***

In the following conditions, the risks of bleeding with Activase are increased and should be weighed against the anticipated benefits: recent major surgery or procedure; cerebrovascular disease; recent intracranial hemorrhage; recent gastrointestinal or genitourinary bleeding; recent trauma; hypertension; acute pericarditis; subacute bacterial endocarditis; hemostatic defects including those secondary to severe hepatic or renal disease; significant hepatic dysfunction; pregnancy; diabetic hemorrhagic retinopathy or other hemorrhagic ophthalmic conditions; septic thrombophlebitis or occluded AV cannula at seriously infected site; advanced age; and patients currently receiving oral anticoagulants, or any other condition in which bleeding constitutes a significant hazard or would be particularly difficult to manage because of its location.

### ***Hypersensitivity***

Hypersensitivity, including urticarial / anaphylactic reactions, have been reported after administration of Activase. Rare fatal outcome for hypersensitivity was reported. Angioedema has been observed during and up to 2 hours after infusion in patients treated for acute ischemic stroke and acute myocardial infarction. In many cases, patients received concomitant angiotensin-converting enzyme inhibitors. Monitor patients during and for several hours after infusion for hypersensitivity. If signs of hypersensitivity occur, e.g. anaphylactoid reaction or angioedema develops, discontinue the Activase infusion and promptly institute appropriate therapy (e.g., antihistamines, intravenous corticosteroids, epinephrine).

12

IMPORTANT SAFETY  
INFORMATION

**ACTIVASE<sup>®</sup>**  
**ALTEPLASE**  
A RECOMBINANT TISSUE PLASMINOGEN ACTIVATOR



# Important Safety Information (*cont'd*)

STROKE ASSESSMENT

TREATMENT

## ***Thromboembolism***

The use of thrombolytics can increase the risk of thrombo-embolic events in patients with high likelihood of left heart thrombus, such as patients with mitral stenosis or atrial fibrillation. Activase has not been shown to treat adequately underlying deep vein thrombosis in patients with PE. Consider the possible risk of re-embolization due to the lysis of underlying deep venous thrombi in this setting.

## ***Cholesterol Embolization***

Cholesterol embolism, sometimes fatal, has been reported rarely in patients treated with thrombolytic agents; the true incidence is unknown. It is associated with invasive vascular procedures (e.g., cardiac catheterization, angiography, vascular surgery) and/or anticoagulant therapy.

## ***Coagulation Tests May be Unreliable during Activase Therapy***

Coagulation tests and/or measures of fibrinolytic activity may be unreliable during Activase therapy unless specific precautions are taken to prevent in vitro artifacts. When present in blood at pharmacologic concentrations, Activase remains active under *in vitro* conditions, which can result in degradation of fibrinogen in blood samples removed for analysis.

## ***Adverse Reactions***

The most frequent adverse reaction associated with Activase AIS therapy is bleeding.

**Please see accompanying full Prescribing Information for additional Important Safety Information.**

12

IMPORTANT SAFETY  
INFORMATION

**ACTIVASE<sup>®</sup>**  
**ALTEPLASE**  
A RECOMBINANT TISSUE PLASMINOGEN ACTIVATOR



# References

## STROKE ASSESSMENT

## TREATMENT

1. Tao W-D, Liu M, Fisher M, Wang D-R, Li J, Furie KL, et al. Posterior versus anterior circulation infarction: how different are the neurological deficits? *Stroke*. 2012;43:2060-2065.
2. Ferbert A, Bruckmann H, Drummen R. Clinical features of proven basilar artery occlusion. *Stroke*. 1990;21:1135-1142.
3. Searls DE, Pazdera L, Korbel E, Vysata O, Caplan LR. Symptoms and signs of posterior circulation ischemia in the New England Medical Center posterior circulation registry. *Arch Neurol*. 2012;69(3):346-351.
4. Jauch EC, French DM, McGeorge T. Prehospital stroke treatment (EMS stabilization protocols). In: Ovbiagele B, Turan TN, eds. *Ischemic Stroke Therapeutics: A Comprehensive Guide*. 1st ed. Switzerland: Springer International; 2016:7-12.
5. NIH Stroke Scale. National Institutes of Health website. [https://www.stroke.nih.gov/documents/NIH\\_Stroke\\_Scale.pdf](https://www.stroke.nih.gov/documents/NIH_Stroke_Scale.pdf). Accessed January 9, 2019.
6. Arch AE, Weisman DC, Coca S, et al. Missed ischemic stroke diagnosis in the emergency department by emergency medicine and neurology services. *Stroke*. 2016;47(3):668-673.
7. Powers WJ, Rabinstein AA, Ackerson T, et al. Guidelines for the early management of patients with acute ischemic stroke: 2019 update to the 2018 guidelines for the early management of acute ischemic stroke: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2019;(12):e344-e418.
8. Demaerschalk BM, Kleindorfer DO, Adeoye OM, et al. Scientific rationale for the inclusion and exclusion criteria for intravenous alteplase in acute ischemic stroke: a statement for healthcare professionals from the American Heart Association/American Stroke Association [published correction appears in *Stroke*. 2016;47:e262]. *Stroke*. 2016;47(2):581-641.
9. Martin-Schild S, Albright KC, Tanksley J, et al. Zero on the NIHSS does not equal the absence of stroke. *Ann Emerg Med*. 2011;57:42-45.
10. NIH Stroke Scale International. Approach to Human Subject Protection and Patient Safety. Available at: <http://www.nihstrokescale.org>. Accessed June 18, 2020.
11. Activase [prescribing information]. South San Francisco, CA: Genentech, Inc; 2018.
12. Winstein CJ, Stein J, Arena R, et al. Guidelines for adult stroke rehabilitation and recovery: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2016;47:e98-e169.

12

## IMPORTANT SAFETY INFORMATION

**ACTIVASE®**  
**ALTEPLASE**  
A RECOMBINANT TISSUE PLASMINOGEN ACTIVATOR



# References (cont'd)

STROKE ASSESSMENT

TREATMENT

- 13.** Banks JL, Marotta CA. Outcomes validity and reliability of the modified Rankin Scale: implications for stroke clinical trials—a literature review and synthesis. *Stroke*. 2007;38:1091-1096. **14.** English JD, Yavagal DR, Gupta R, et al. Mechanical thrombectomy-ready comprehensive stroke center requirements and endovascular stroke systems of care: recommendations from the endovascular stroke standards committee of the Society of Vascular and Interventional Neurology (SVIN). *Intervent Neurol*. 2015;4:138-150. **15.** The Joint Commission. Prepublication requirements. Requirement for primary stroke center advanced certification. Issued January 17, 2014. **16.** American Heart Association/American Stroke Association. *Target: Stroke Phase II Campaign Manual*. Published October 2014. **17.** Summers D, Leonard A, Wentworth D, et al. Comprehensive overview of nursing and interdisciplinary care of the acute ischemic stroke patient: a scientific statement from the American Heart Association. *Stroke*. 2009;40:2911-2944.

12

IMPORTANT SAFETY  
INFORMATION

**ACTIVASE®**  
**ALTEPLASE**  
A RECOMBINANT TISSUE PLASMINOGEN ACTIVATOR







**Genentech**

*A Member of the Roche Group*

© 2020 Genentech USA, Inc.

All rights reserved. M-US-00006249(v1.0)

**ACTIVASE<sup>®</sup>**  
**ALTEPLASE**

A RECOMBINANT TISSUE PLASMINOGEN ACTIVATOR



## HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use ACTIVASE safely and effectively. See full prescribing information for ACTIVASE.

Activase (alteplase) for injection, for intravenous use

Initial U.S. Approval: 1987

### -----RECENT MAJOR CHANGES-----

Warnings and Precautions (5.3) 02/2018

### -----INDICATIONS AND USAGE-----

Activase is a tissue plasminogen activator (tPA) indicated for the treatment of

- Acute Ischemic Stroke (AIS). (1.1)
- Acute Myocardial Infarction (AMI) to reduce mortality and incidence of heart failure. (1.2)

Limitation of Use in AMI: the risk of stroke may be greater than the benefit in patients at low risk of death from cardiac causes. (1.2)

- Acute Massive Pulmonary Embolism (PE) for lysis. (1.3)

### -----DOSAGE AND ADMINISTRATION-----

- Acute Ischemic Stroke: The recommended dose is 0.9 mg/kg (not to exceed 90 mg total dose) infused intravenously over 60 minutes with 10% of the total dose administered as an initial bolus over 1 minute. (2.1)
- Acute Myocardial Infarction: The recommended total dose is based on patient weight, not to exceed 100 mg. (2.2)
- Acute Massive Pulmonary Embolism: The recommended dose is 100 mg administered by IV infusion over 2 hours. (2.3)
- Do not add other medications to infusions containing Activase. (2.5)

### -----DOSAGE FORMS AND STRENGTHS-----

- Lyophilized powder: 50 mg and 100 mg with Sterile Water for Injection USP for reconstitution at 1 mg per 1 mL. (3)

### -----CONTRAINDICATIONS-----

#### General

- Active internal bleeding. (4.1, 4.2)
- Recent intracranial or intraspinal surgery or serious head trauma. (4.1, 4.2)
- Intracranial conditions that may increase the risk of bleeding. (4.1, 4.2)
- Bleeding diathesis. (4.1, 4.2)
- Current severe uncontrolled hypertension. (4.1, 4.2)

#### Acute Ischemic Stroke

- Current intracranial hemorrhage. (4.1)
- Subarachnoid hemorrhage. (4.1)

#### Acute Myocardial Infarction or Pulmonary Embolism

- History of recent stroke. (4.2)

### -----WARNINGS AND PRECAUTIONS-----

- Increases the risk of bleeding. Avoid intramuscular injections. Monitor for bleeding. If serious bleeding occurs, discontinue Activase. (5.1)
- Monitor patients during and for several hours after infusion for hypersensitivity. If signs of hypersensitivity develop, discontinue Activase. (5.2)
- Consider the risk of reembolization from the lysis of underlying deep venous thrombi in patients with pulmonary embolism. (5.3)
- Cholesterol embolism has been reported rarely in patients treated with thrombolytic agents. (5.4)

### -----ADVERSE REACTIONS-----

The most frequently occurring adverse reaction (> 5%) is bleeding. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Genentech at 1-888-835-2555 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

### -----DRUG INTERACTIONS-----

- Anticoagulants and drugs that inhibit platelet function increase the risk of bleeding when administered with Activase therapy. (7)
- Concomitant angiotensin-converting enzyme inhibitors may increase the risk of angioedema. (7)

### -----USE IN SPECIFIC POPULATIONS-----

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 02/2018

## FULL PRESCRIBING INFORMATION: CONTENTS\*

### 1 INDICATIONS AND USAGE

- 1.1 Acute Ischemic Stroke
- 1.2 Acute Myocardial Infarction
- 1.3 Pulmonary Embolism

### 2 DOSAGE AND ADMINISTRATION

- 2.1 Acute Ischemic Stroke
- 2.2 Acute Myocardial Infarction
- 2.3 Pulmonary Embolism
- 2.4 Preparation for Administration
- 2.5 Administration

### 3 DOSAGE FORMS AND STRENGTHS

### 4 CONTRAINDICATIONS

- 4.1 Acute Ischemic Stroke
- 4.2 Acute Myocardial Infarction or Pulmonary Embolism

### 5 WARNINGS AND PRECAUTIONS

- 5.1 Bleeding
- 5.2 Hypersensitivity
- 5.3 Thromboembolism
- 5.4 Cholesterol Embolization
- 5.5 Coagulation Tests May Be Unreliable during Activase Therapy

### 6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Post-Marketing Experience

### 7 DRUG INTERACTIONS

### 8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use

### 11 DESCRIPTION

### 12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

### 13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

### 14 CLINICAL STUDIES

- 14.1 Acute Ischemic Stroke (AIS)
- 14.2 Acute Myocardial Infarction (AMI)
- 14.3 Acute Massive Pulmonary Embolism (PE)

### 16 HOW SUPPLIED/STORAGE AND HANDLING

- 16.1 How Supplied
- 16.2 Stability and Storage

### 17 PATIENT COUNSELING INFORMATION

\* Sections or subsections omitted from the Full Prescribing Information are not listed.

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

#### **1.1 Acute Ischemic Stroke**

Activase is indicated for the treatment of acute ischemic stroke.

Exclude intracranial hemorrhage as the primary cause of stroke signs and symptoms prior to initiation of treatment [see *Contraindications (4.1)*]. Initiate treatment as soon as possible but within 3 hours after symptom onset.

#### **1.2 Acute Myocardial Infarction**

Activase is indicated for use in acute myocardial infarction (AMI) for the reduction of mortality and reduction of the incidence of heart failure.

Limitation of Use: The risk of stroke may outweigh the benefit produced by thrombolytic therapy in patients whose AMI puts them at low risk for death or heart failure.

#### **1.3 Pulmonary Embolism**

Activase is indicated for the lysis of acute massive pulmonary embolism, defined as:

- Acute pulmonary emboli obstructing blood flow to a lobe or multiple lung segments.
- Acute pulmonary emboli accompanied by unstable hemodynamics, e.g., failure to maintain blood pressure without supportive measures.

### **2 DOSAGE AND ADMINISTRATION**

#### **2.1 Acute Ischemic Stroke**

Administer Activase as soon as possible but within 3 hours after onset of symptoms.

The recommended dose is 0.9 mg/kg (not to exceed 90 mg total dose), with 10% of the total dose administered as an initial intravenous bolus over 1 minute and the remainder infused over 60 minutes.

During and following Activase administration for the treatment of acute ischemic stroke, frequently monitor and control blood pressure.

In patients without recent use of oral anticoagulants or heparin, Activase treatment can be initiated prior to the availability of coagulation study results. Discontinue Activase if the pretreatment International Normalized Ratio (INR) is greater than 1.7 or the activated partial thromboplastin time (aPTT) is elevated [see *Contraindications (4.1)*].

#### **2.2 Acute Myocardial Infarction**

Administer Activase as soon as possible after the onset of symptoms.

The recommended total doses for acute myocardial infarction (AMI) is based on patient weight, not to exceed 100 mg, regardless of the selected administration regimen (accelerated or 3 hour, described below).

There are two Activase dose regimens (accelerated and 3-hour) for use in the management of AMI; there are no controlled studies to compare clinical outcomes with these regimens [see *Clinical Studies (14.2)*].

### Accelerated Infusion

The recommended accelerated infusion dose consists of an IV bolus [see *Dosage and Administration (2.5)*] followed by an IV infusion as set forth in Table 1.

**Table 1**  
Accelerated Infusion Weight-Based Doses for Patients with AMI

Patient weight	Intravenous Bolus	First 30 min	Next 60 min
> 67 kg	15 mg	50 mg	35 mg
≤ 67 kg	15 mg	0.75 mg/kg	0.50 mg/kg

The safety and efficacy of accelerated infusion of Activase have only been investigated with concomitant administration of heparin and aspirin [see *Clinical Studies (14.2)*].

### 3-Hour Infusion

For patients weighing  $\geq 65$  kg, the recommended dose is 100 mg administered as 60 mg in the first hour (6-10 mg administered as a bolus), 20 mg over the second hour, and 20 mg over the third hour. For smaller patients (< 65 kg), a dose of 1.25 mg/kg administered over 3 hours may be used. Weight-based doses are shown in Table 2.

**Table 2**  
3-hour Infusion Weight-Based Doses for Patients with AMI

Patient weight	Bolus	Rest of 1st hour	2nd hour	3rd hour
$\geq 65$ kg	6-10 mg	50-54 mg	20 mg	20 mg
< 65 kg	0.075 mg/kg	0.675 mg/kg	0.25 mg/kg	0.25 mg/kg

## 2.3 Pulmonary Embolism (PE)

The recommended dose is 100 mg administered by IV infusion over 2 hours.

Institute parenteral anticoagulation near the end of or immediately following the Activase infusion when the partial thromboplastin time or thrombin time returns to twice normal or less.

## 2.4 Preparation for Administration

### Reconstitution

Use only the accompanying Sterile Water for Injection (SWFI), USP without preservatives. Do not use Bacteriostatic Water for Injection, USP.

Reconstitute using aseptic technique. Do not add other medication to solutions containing Activase. Reconstitute Activase no more than 8 hours before use, as it contains no antibacterial preservatives [see *How Supplied/Storage and Handling (16.2)*].

Slight foaming is not unusual; let stand undisturbed for several minutes to allow large bubbles to dissipate. Inspect parenteral drug products for particulate matter and discoloration prior to administration whenever solution and container permit.

Activase may be administered as reconstituted at 1 mg/mL or further diluted immediately before administration in an equal volume of 0.9% Sodium Chloride Injection, USP, or 5% Dextrose Injection, USP, to yield a concentration of 0.5 mg/mL, using either polyvinyl chloride bags or glass vials.

Avoid excessive agitation during dilution; mix by gently swirling and/or slow inversion.

### 50 mg Vials

DO NOT USE IF VACUUM IS NOT PRESENT.

Using a large bore needle (e.g., 18 gauge) and a syringe, reconstitute by adding the contents of the accompanying 50 mL vial of SWFI to the 50 mg vial of Activase, directing the SWFI stream into the lyophilized cake.

### 100 mg Vials

THE 100 mg VIALS DO NOT CONTAIN VACUUM.

Using the transfer device provided, reconstitute by adding the contents of the accompanying 100 mL vial of SWFI to the 100 mg vial of Activase.

1. Use aseptic technique.
2. Remove the protective flip-caps from one vial of Activase and one vial of SWFI.
3. Open the package containing the transfer device by peeling the paper label off the package.
4. Remove the protective cap from one end of the transfer device and keeping the vial of SWFI upright, insert the piercing pin vertically into the center of the stopper of the vial of SWFI.
5. Remove the protective cap from the other end of the transfer device. DO NOT INVERT THE VIAL OF SWFI.
6. Hold the vial of Activase upside down, position it so that the center of the stopper is directly over the exposed piercing pin of the transfer device, and push the vial of Activase down so that the piercing pin is inserted through the center of the Activase vial stopper.
7. Invert the two vials so that the vial of Activase is on the bottom (upright) and the vial of SWFI is upside-down, allowing the SWFI to flow down through the transfer device. Allow the entire contents of the vial of SWFI to flow into the Activase vial (approximately 0.5 cc of SWFI will remain in the diluent vial).
8. Remove the transfer device and the empty SWFI vial from the Activase vial and discard.
9. Swirl gently to dissolve the Activase powder. DO NOT SHAKE.

### **Preparation of Bolus Dose**

- Prepare the bolus dose in one of the following ways: Remove the appropriate volume from the vial of reconstituted (1 mg/mL) Activase using a syringe and needle. If this method is used with the 50 mg vials, the syringe should not be primed with air and the needle should be inserted into the Activase vial stopper. If the 100 mg vial is used, the needle should be inserted away from the puncture mark made by the transfer device.
- Remove the appropriate volume from a port (second injection site) on the infusion line after the infusion set is primed.

- Program an infusion pump to deliver the appropriate volume as a bolus at the initiation of the infusion

## **2.5 Administration**

Following bolus dose, if indicated [*see Dosage and Administration (2.1, 2.2)*]:

- 50 mg vials - administer using either a polyvinyl chloride bag or glass vial and infusion set.
- 100 mg vials - remove from the vial any quantity of drug in excess of that specified for patient treatment [*see Dosage and Administration (2.1, 2.2)*]. Insert the spike end of an infusion set through the same puncture site created by the transfer device in the stopper of the vial of reconstituted Activase. Peel the clear plastic hanger from the vial label. Hang the Activase vial from the resulting loop.

Activase is for intravenous administration only. Extravasation of Activase infusion can cause ecchymosis or inflammation. If extravasation occurs, terminate the infusion at that IV site and apply local therapy.

Do not add any other medication to infusion solutions containing Activase.

## **3 DOSAGE FORMS AND STRENGTHS**

- 50 mg lyophilized powder per single use vial with 50 mL SWFI USP for reconstitution
- 100 mg lyophilized powder per single use vial with 100 mL SWFI USP for reconstitution

## **4 CONTRAINDICATIONS**

### **4.1 Acute Ischemic Stroke**

Do not administer Activase to treat acute ischemic stroke in the following situations in which the risk of bleeding is greater than the potential benefit [*see Warnings and Precautions (5.1)*]:

- Current intracranial hemorrhage
- Subarachnoid hemorrhage
- Active internal bleeding
- Recent (within 3 months) intracranial or intraspinal surgery or serious head trauma
- Presence of intracranial conditions that may increase the risk of bleeding (e.g., some neoplasms, arteriovenous malformations, or aneurysms)
- Bleeding diathesis
- Current severe uncontrolled hypertension.

### **4.2 Acute Myocardial Infarction or Pulmonary Embolism**

Do not administer Activase for treatment of AMI or PE in the following situations in which the risk of bleeding is greater than the potential benefit [*see Warnings and Precautions (5.1)*]:

- Active internal bleeding
- History of recent stroke
- Recent (within 3 months) intracranial or intraspinal surgery or serious head trauma
- Presence of intracranial conditions that may increase the risk of bleeding (e.g. some neoplasms, arteriovenous malformations, or aneurysms)

- Bleeding diathesis
- Current severe uncontrolled hypertension.

## **5 WARNINGS AND PRECAUTIONS**

### **5.1 Bleeding**

Activase can cause significant, sometimes fatal, internal or external bleeding, especially at arterial and venous puncture sites. Avoid intramuscular injections and trauma to the patient while on Activase. Perform venipunctures carefully and only as required. To minimize bleeding from noncompressible sites, avoid internal jugular and subclavian venous punctures. If an arterial puncture is necessary during Activase infusion, use an upper extremity vessel that is accessible to manual compression, apply pressure for at least 30 minutes, and monitor the puncture site closely.

Because of the higher risk of intracranial hemorrhage in patients treated for acute ischemic stroke, limit treatment to facilities that can provide timely access to appropriate evaluation and management of intracranial hemorrhage.

Fatal cases of hemorrhage associated with traumatic intubation in patients administered Activase have been reported.

Aspirin and heparin have been administered concomitantly with and following infusions of Activase in the management of acute myocardial infarction and pulmonary embolism, but the concomitant administration of heparin and aspirin with and following infusions of Activase for the treatment of acute ischemic stroke during the first 24 hours after symptom onset has not been investigated. Because heparin, aspirin, or Activase may cause bleeding complications, carefully monitor for bleeding, especially at arterial puncture sites. Hemorrhage can occur 1 or more days after administration of Activase, while patients are still receiving anticoagulant therapy.

If serious bleeding occurs, terminate the Activase infusion and treat appropriately. In the following conditions, the risks of bleeding with Activase therapy for all approved indications are increased and should be weighed against the anticipated benefits:

- Recent major surgery or procedure, (e.g., coronary artery bypass graft, obstetrical delivery, organ biopsy, previous puncture of noncompressible vessels)
- Cerebrovascular disease
- Recent intracranial hemorrhage
- Recent gastrointestinal or genitourinary bleeding
- Recent trauma
- Hypertension: systolic BP above 175 mm Hg or diastolic BP above 110 mm Hg
- Acute pericarditis
- Subacute bacterial endocarditis
- Hemostatic defects including those secondary to severe hepatic or renal disease
- Significant hepatic dysfunction
- Pregnancy
- Diabetic hemorrhagic retinopathy, or other hemorrhagic ophthalmic conditions
- Septic thrombophlebitis or occluded AV cannula at seriously infected site

Advanced age [*see Use in Specific Populations (8.5)*]

- Patients currently receiving anticoagulants (e.g., warfarin sodium)

Any other condition in which bleeding constitutes a significant hazard or would be particularly difficult to manage because of its location.

## **5.2 Hypersensitivity**

Hypersensitivity, including urticarial / anaphylactic reactions, have been reported after administration of Activase (e.g., laryngeal edema, rash and shock). Rare fatal outcome for hypersensitivity was reported. Angioedema has been observed during and up to 2 hours after Activase infusion in patients treated for acute ischemic stroke and acute myocardial infarction. In many cases, patients received concomitant angiotensin-converting enzyme inhibitors [*see Drug Interactions (7)*].

Monitor patients treated with Activase during and for several hours after infusion for hypersensitivity. If signs of hypersensitivity occur, e.g. anaphylactoid reaction or angioedema develops, discontinue the Activase infusion and promptly institute appropriate therapy (e.g., antihistamines, intravenous corticosteroids, epinephrine).

## **5.3 Thromboembolism**

The use of thrombolytics can increase the risk of thrombo-embolic events in patients with high likelihood of left heart thrombus, such as patients with mitral stenosis or atrial fibrillation. Activase has not been shown to treat adequately underlying deep vein thrombosis in patients with PE. Consider the possible risk of re-embolization due to the lysis of underlying deep venous thrombi in this setting.

## **5.4 Cholesterol Embolization**

Cholesterol embolism has been reported rarely in patients treated with thrombolytic agents; the true incidence is unknown. Cholesterol embolism may present with livedo reticularis, “purple toe” syndrome, acute renal failure, gangrenous digits, hypertension, pancreatitis, myocardial infarction, cerebral infarction, spinal cord infarction, retinal artery occlusion, bowel infarction, or rhabdomyolysis and can be fatal. It is associated with invasive vascular procedures (e.g., cardiac catheterization, angiography, vascular surgery) and/or anticoagulant therapy.

## **5.5 Coagulation Tests May Be Unreliable during Activase Therapy**

Coagulation tests and measures of fibrinolytic activity may be unreliable during Activase therapy, unless specific precautions are taken to prevent in vitro artifacts. When present in blood at pharmacologic concentrations, Activase remains active under in vitro conditions, which can result in degradation of fibrinogen in blood samples removed for analysis.

## **6 ADVERSE REACTIONS**

The following adverse reactions are discussed in greater detail in the other sections of the label:

- Bleeding [*see Contraindications (4), Warnings and Precautions (5.1)*]
- Hypersensitivity [*see Warnings and Precautions (5.2)*]
- Thromboembolism [*see Warnings and Precautions (5.3)*]
- Cholesterol Embolization [*see Warnings and Precautions (5.4)*]

## 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

The most frequent adverse reaction associated with Activase in all approved indications is bleeding.

### ***Bleeding***

#### *Acute Ischemic Stroke (AIS)*

In clinical studies in patients with AIS (Studies 1 and 2) the incidence of intracranial hemorrhage, especially symptomatic intracranial hemorrhage, was higher in Activase-treated patients than in placebo patients. A dose-finding study of Activase suggested that doses greater than 0.9 mg/kg may be associated with an increased incidence of intracranial hemorrhage.

The incidence of all-cause 90-day mortality, intracranial hemorrhage, and new ischemic stroke following Activase treatment compared to placebo are presented in Table 3 as a combined safety analysis (n=624) for Studies 1 and 2. These data indicate a significant increase in intracranial hemorrhage following Activase treatment, particularly symptomatic intracranial hemorrhage within 36 hours. There was no increase in the incidences of 90-day mortality or severe disability in Activase-treated patients compared to placebo.

**Table 3**  
Combined Safety Outcomes for Studies 1 and 2

	Placebo (n=312)	Activase (n=312)	p-Value <sup>b</sup>
All-Cause 90-day Mortality	64 (20.5%)	54 (17.3%)	0.36
Total ICH <sup>a</sup>	20 (6.4%)	48 (15.4%)	<0.01
Symptomatic	4 (1.3%)	25 (8.0%)	<0.01
Asymptomatic	16 (5.1%)	23 (7.4%)	0.32
Symptomatic Intracranial Hemorrhage within 36 hours	2 (0.6%)	20 (6.4%)	<0.01
New Ischemic Stroke (3-months)	17 (5.4%)	18 (5.8%)	1.00

<sup>a</sup> Within trial follow-up period. Symptomatic intracranial hemorrhage was defined as the occurrence of sudden clinical worsening followed by subsequent verification of intracranial hemorrhage on CT scan. Asymptomatic intracranial hemorrhage was defined as intracranial hemorrhage detected on a routine repeat CT scan without preceding clinical worsening.

<sup>b</sup> Fisher's Exact Test.



Bleeding events other than intracranial hemorrhage were noted in the studies of AIS and were consistent with the general safety profile of Activase. In Studies 1 and 2, the frequency of bleeding requiring red blood cell transfusions was 6.4% for Activase-treated patients compared to 3.8% for placebo (p=0.19).

Although exploratory analyses of Studies 1 and 2 suggest that severe neurological deficit (National Institutes of Health Stroke Scale [NIHSS > 22]) at presentation was associated with an increased risk of intracranial hemorrhage, efficacy results suggest a reduced but still favorable clinical outcome for these patients.

#### *Acute Myocardial Infarction (AMI)*

For the 3-hour infusion regimen in the treatment of AMI, the incidence of significant internal bleeding (estimated as > 250 mL blood loss) has been reported in studies in over 800 patients (Table 4). These data do not include patients treated with the Activase accelerated infusion.

**Table 4**  
Incidence of Bleeding in 3-Hour Infusion in AMI Patients

	Total Dose ≤100 mg
Gastrointestinal	5%
Genitourinary	4%
Ecchymosis	1%
Retroperitoneal	<1%
Epistaxis	<1%
Gingival	<1%

The incidence of intracranial hemorrhage in AMI patients treated with Activase is presented in Table 5.

**Table 5**  
Incidence of Intracranial Hemorrhage in AMI Patients

Dose	Number of Patients	Intracranial Hemorrhage (%)
100 mg, 3-hour	3272	0.4
≤ 100 mg, accelerated	10,396	0.7
150 mg	1779	1.3
1-1.4 mg/kg	237	0.4

A dose of 150 mg or greater should not be used in the treatment of AMI because it has been associated with an increase in intracranial bleeding.

#### *Pulmonary Embolism (PE)*

For acute massive pulmonary embolism, bleeding events were consistent with the general safety profile observed with Activase treatment of AMI patients receiving the 3-hour infusion regimen.

## 6.2 Post-Marketing Experience

The following adverse reactions have been identified during post-approval use of Activase. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure. These reactions are frequent sequelae of the underlying disease, and the effect of Activase on the incidence of these events is unknown.

*Acute Ischemic Stroke:* Cerebral edema, cerebral herniation, seizure, new ischemic stroke, embolism. These events may be life threatening and may lead to death.

*Acute Myocardial Infarction:* Arrhythmias, AV block, cardiogenic shock, heart failure, cardiac arrest, recurrent ischemia, myocardial reinfarction, myocardial rupture, electromechanical dissociation, pericardial effusion, pericarditis, mitral regurgitation, cardiac tamponade, thromboembolism, pulmonary edema. These events may be life threatening and may lead to death. Nausea and/or vomiting, hypotension and fever have also been reported.

*Pulmonary Embolism:* Pulmonary reembolization, pulmonary edema, pleural effusion, thromboembolism, hypotension. These events may be life threatening and may lead to death. Fever has also been reported.

## 7 DRUG INTERACTIONS

The interaction of Activase with other cardioactive or cerebroactive drugs has not been studied. Anticoagulants and antiplatelet drugs increase the risk of bleeding if administered prior to, during, or after Activase therapy.

In the post-marketing setting, there have been reports of angioedema in patients (primarily patients with AIS) receiving concomitant angiotensin-converting enzyme inhibitors. [see *Warnings and Precautions* (5.2)].

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

#### Risk Summary

Published studies and case reports on alteplase use in pregnant women are insufficient to inform a drug associated risk of adverse developmental outcomes. Alteplase is embryocidal in rabbits when intravenously administered during organogenesis at the clinical exposure for AMI, but no maternal or fetal toxicity was evident at lower exposure in pregnant rats or rabbits (*see Data*).

All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. The estimated background risk of major birth defects and miscarriage for the indicated populations is unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2-4% and 15-20%, respectively.

#### Clinical Considerations

##### *Maternal Adverse Reactions*

The most common complication of thrombolytic therapy is bleeding. Pregnancy may increase this risk [see *Warnings and Precautions* (5.1)].

#### Data

## *Animal Data*

Alteplase is embryocidal in rabbits when administered intravenously during organogenesis in doses (3 mg/kg) approximately equal to the human exposure (based on AUC) at the dose for AMI. No maternal or fetal toxicity was evident at doses (1 mg/kg) approximately 0.3 times the human exposure. In pregnant rats, no maternal or fetal toxicity was evident at doses (1 mg/kg) approximately 0.6 times the human dose for AMI (based on body weight) dosed during the period of organogenesis.

## **8.2 Lactation**

### Risk Summary

There are no data on the presence of alteplase in human milk, the effects on the breastfed infant, or the effects on milk production.

## **8.4 Pediatric Use**

Safety and effectiveness of Activase in pediatric patients have not been established.

## **8.5 Geriatric Use**

### *Acute Ischemic Stroke*

In exploratory, multivariate analyses of Studies 1 and 2, age greater than 77 years was one of several interrelated baseline characteristics associated with an increased risk of intracranial hemorrhage. Efficacy results suggest a reduced but still favorable clinical outcome for Activase-treated elderly [see *Clinical Studies (14.1)*].

### *Acute Myocardial Infarction*

In a large trial of accelerated-infusion Activase that enrolled 41,021 patients with AMI to one of four thrombolytic regimens [see *Clinical Studies (14.2)*], patients over 75 years of age, a predefined subgroup, comprised 12% of enrollment. In these patients, the incidence of stroke was 4.0% for the Activase accelerated infusion group, 2.8% for streptokinase IV [SK (IV)], and 3.2% for streptokinase SQ [SK (SQ)]. The incidence of combined 30-day mortality or nonfatal stroke was 20.6% for accelerated infusion of Activase, 21.5% for SK (IV), and 22.0% for SK (SQ).

## **11 DESCRIPTION**

Activase is a tissue plasminogen activator produced by recombinant DNA technology. It is a sterile, purified glycoprotein of 527 amino acids. It is synthesized using the complementary DNA (cDNA) for natural human tissue-type plasminogen activator obtained from a human melanoma cell line. Activase is a sterile, white to off-white, lyophilized powder for intravenous administration after reconstitution with Sterile Water for Injection, USP.

**Table 6**  
Quantitative Composition of the Lyophilized Product

	100 mg Vial	50 mg Vial
Alteplase	100 mg (58 million IU)	50 mg (29 million IU)
L-Arginine	3.5 g	1.7 g
Phosphoric Acid	1 g	0.5 g
Polysorbate 80	10 mg	5 mg
Vacuum	No	Yes

Biological potency is determined by an in vitro clot lysis assay and is expressed in International Units (IU).

The reconstituted preparation results in a colorless to pale yellow transparent solution containing Activase 1 mg/mL at approximately pH 7.3. The osmolality of this solution is approximately 215 mOsm/kg.

## **12 CLINICAL PHARMACOLOGY**

### **12.1 Mechanism of Action**

Alteplase is a serine protease responsible for fibrin-enhanced conversion of plasminogen to plasmin. It produces limited conversion of plasminogen in the absence of fibrin.

When introduced into the systemic circulation at pharmacologic concentration, alteplase binds to fibrin in a thrombus and converts the entrapped plasminogen to plasmin. This initiates local fibrinolysis with limited systemic proteolysis.

### **12.2 Pharmacodynamics**

Following administration of 100 mg Activase, there is a decrease (16%-36%) in circulating fibrinogen. In a controlled trial, 8 of 73 patients (11%) receiving Activase (1.25 mg/kg body weight over 3 hours) experienced a decrease in fibrinogen to below 100 mg/dL.

### **12.3 Pharmacokinetics**

Alteplase in acute myocardial infarction (AMI) patients is rapidly cleared from the plasma with an initial half-life of less than 5 minutes. There is no difference in the dominant initial plasma half-life between the 3-hour and accelerated regimens for AMI. The plasma clearance of alteplase is 380-570 mL/min, primarily mediated by the liver. The initial volume of distribution approximates plasma volume.

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

Long-term studies in animals have not been performed to evaluate the carcinogenic potential or the effect on fertility. Short-term studies, which evaluated tumorigenicity of Activase and effect on tumor metastases in rodents, were negative.

Studies to determine mutagenicity (Ames test) and chromosomal aberration assays in human lymphocytes were negative at all concentrations tested. Cytotoxicity, as reflected by a decrease in mitotic index, was evidenced only after prolonged exposure and only at the highest concentrations tested.

## **14 CLINICAL STUDIES**

### **14.1 Acute Ischemic Stroke (AIS)**

Two placebo-controlled, double-blind trials (Studies 1 and 2) were conducted in patients with AIS. Both studies enrolled patients with measurable neurological deficit who could complete screening and begin study treatment within 3 hours from symptom onset. A cranial computerized tomography (CT) scan was performed prior to treatment to rule out the presence of intracranial hemorrhage. Blood pressure was actively controlled (185/110 mm Hg or lower) for 24 hours.

Patients were randomized (1:1) to receive either 0.9 mg/kg Activase (maximum of 90 mg) or placebo. Activase was administered as a 10% initial IV bolus over 1 minute followed by continuous IV infusion of the remainder over 60 minutes. Study treatment was initiated prior to the availability of coagulation study results in patients without recent use of oral anticoagulants and/or heparin and was discontinued if the pretreatment prothrombin time (PT) was greater than 15 seconds or the activated partial thromboplastin time (aPTT) was elevated. Patients with prior aspirin use were included. Administration of anticoagulants and antiplatelet agents was prohibited for the first 24 hours following symptom onset.

Study 1 (n=291) evaluated neurological improvement at 24 hours after stroke onset. The primary endpoint, the proportion of patients with a 4 point or greater improvement in the National Institutes of Health Stroke Scale (NIHSS) score or complete recovery (NIHSS score of 0), was not significantly different between treatment groups. A prespecified secondary analysis suggested improved 3-month outcome associated with Activase treatment using the following stroke assessment scales: Barthel Index, Modified Rankin Scale, Glasgow Outcome Scale, and the NIHSS.

Study 2 (n=333) assessed clinical outcome at 3 months. A favorable outcome was defined as minimal or no disability using four stroke assessment scales: Barthel Index (score of 95 or greater), Modified Rankin Scale (score of 1 or less), Glasgow Outcome Scale (score of 1), and NIHSS (score of 1 or less). The results comparing Activase- and placebo-treated patients for the four outcome scales together (Generalized Estimating Equations) and individually are presented in Table 7. In this study, depending upon the scale, the favorable outcome of minimal or no disability occurred in at least 11 per 100 more patients treated with Activase than those receiving placebo. Study results demonstrated consistent functional and neurological improvement within all four stroke scales as indicated by median scores. These results were consistent with the 3-month outcome treatment effects observed in Study 1.

**Table 7**  
Study 2 Three-Month Efficacy Outcomes

Analysis	Frequency of Favorable Outcome <sup>a</sup>				p-Value <sup>c</sup>
	Placebo (n=165)	Activase (n=168)	Absolute Difference (95% CI)	Odds Ratio <sup>b</sup> (95% CI)	
Generalized Estimating Equations (Multivariate)	-	-	-	1.71 (1.15, 2.56)	0.02
Barthel Index	37.6%	50.0%	12.4% (3.0, 21.9)	1.66 (1.07, 2.57)	0.02
Modified Rankin Scale	26.1%	38.7%	12.6% (3.7, 21.6)	1.79 (1.12, 2.85)	0.02
Glasgow Outcome Scale	31.5%	44.0%	12.5% (3.3, 21.8)	1.71 (1.09, 2.68)	0.02
NIHSS	20.0%	31.0%	11.0% (2.6, 19.3)	1.79 (1.06, 2.96)	0.02

<sup>a</sup> Favorable Outcome is defined as recovery with minimal or no disability.

<sup>b</sup> Value greater than 1 indicates odds of recovery in favor of Activase treatment.

<sup>c</sup> p-Value for Odds Ratio is from Generalized Estimating Equations with logit link.

In a prespecified subgroup analysis of patients receiving aspirin prior to onset of stroke symptoms, the favorable outcome for Activase-treated patients was preserved.

## 14.2 Acute Myocardial Infarction (AMI)

Two Activase dose regimens have been studied in patients experiencing acute myocardial infarction [see *Dosage and Administration* (2.2)]. The comparative efficacy of these two regimens has not been evaluated.

### *Accelerated Infusion in AMI Patients*

Accelerated infusion of Activase was studied in an international, multi-center trial that randomized 41,021 patients with AMI to four thrombolytic regimens (Study 3). Entry criteria included onset of chest pain within 6 hours of treatment and ST-segment elevation of ECG. The four treatment regimens included accelerated infusion of Activase ( $\leq 100$  mg over 90 minutes) plus intravenous (IV) heparin (n=10,396); Streptokinase (1.5 million units over 60 minutes) plus IV heparin (SK [IV], n=10,410); Streptokinase plus subcutaneous (SQ) heparin (SK [SQ] n=9841). A fourth regimen combined Activase and Streptokinase (n=10,374). All patients received 160 mg chewable aspirin administered as soon as possible, followed by 160-325 mg daily. Bolus IV heparin 5000 U was initiated as soon as possible, followed by a 1000 U/hour continuous IV infusion for at least 48 hours; subsequent heparin therapy was at the physician's discretion. Heparin SQ 12,500 U was administered 4 hours after initiation of SK therapy, followed by 12,500 U twice daily for 7 days or until discharge, whichever came first. Many of the patients randomized to receive SQ heparin

received some IV heparin, usually in response to recurrent chest pain and/or the need for a medical procedure. Some received IV heparin on arrival to the emergency room prior to enrollment and randomization.

Key results from Study 3 are shown in Table 8. The incidence of 30-day mortality for Activase accelerated infusion was 1.0% lower than for either Streptokinase plus heparin regimen. The incidence of combined 30-day mortality or nonfatal stroke for the Activase accelerated infusion was 1.0% lower than for SK (IV) and 0.8% lower than for SK (SQ).

**Table 8**  
Efficacy and Safety Results for Study 3

Event	Accelerated Activase	SK (IV)	p-Value <sup>a</sup>	SK (SQ)	p-Value <sup>a</sup>
30-Day Mortality	6.3%	7.3%	0.003	7.3%	0.007
30-Day Mortality or Nonfatal Stroke	7.2%	8.2%	0.006	8.0%	0.036
24-Hour Mortality	2.4%	2.9%	0.009	2.8%	0.029
Any Stroke	1.6%	1.4%	0.32	1.2%	0.03
Intracerebral Hemorrhage	0.7%	0.6%	0.22	0.5%	0.02

<sup>a</sup> Two-tailed p-value is for comparison of Accelerated Activase to the respective SK control arm.

Subgroup analysis of patients by age, infarct location, time from symptom onset to thrombolytic treatment, and treatment in the U.S. or elsewhere showed consistently lower 30-day mortality on Activase.

For patients who were over 75 years of age, a predefined subgroup consisting of 12% of patients enrolled, the incidence of stroke was 4.0% for the Activase accelerated infusion group, 2.8% for SK (IV), and 3.2% for SK (SQ); the incidence of combined 30-day mortality or nonfatal stroke was 20.6% for accelerated infusion of Activase, 21.5% for SK (IV), and 22.0% for SK (SQ).

### *3-Hour Infusion in AMI Patients*

In a double-blind, randomized trial (n=138) comparing 3-hour infusion of Activase to placebo (Study 4), patients infused with Activase within 4 hours of onset of symptoms experienced improved left ventricular function at Day 10 compared to the placebo group, when ejection fraction was measured by gated blood pool scan (53.2% vs. 46.4%, p=0.018). Relative to baseline (Day 1) values, the net changes in ejection fraction were +3.6% and -4.7% for the treated and placebo groups, respectively (p=0.0001). The treated group had a reduced incidence of clinical heart failure (14%) compared to the placebo group (33%) (p=0.009).

In a double-blind, randomized trial (n=5013) comparing 3-hour infusion of Activase to placebo (Study 5), patients infused with Activase within 5 hours of AMI symptom onset experienced improved 30-day survival compared to the placebo arm. At 1 month, the overall mortality rates were 7.2% for the Activase group and 9.8% for the placebo group (p=0.001). At 6 months, the overall mortality rate for Activase-treated patients was 10.4% compared to the placebo arm (13.1%, p=0.008).

### 14.3 Acute Massive Pulmonary Embolism (PE)

Study 6 was a comparative randomized trial (n=45) in which 59% of patients (n=22) treated with Activase (100 mg over 2 hours) experienced moderate or marked lysis of pulmonary emboli when assessed by pulmonary angiography 2 hours after treatment initiation. Activase-treated patients also experienced a significant reduction in pulmonary embolism-induced pulmonary hypertension within 2 hours of treatment (p=0.003). Pulmonary perfusion at 24 hours, as assessed by radionuclide scan, was significantly improved (p=0.002).

## 16 HOW SUPPLIED/STORAGE AND HANDLING

### 16.1 How Supplied

Activase is supplied as a sterile, lyophilized powder in 50 mg vials containing vacuum and in 100 mg vials without vacuum.

Each 50 mg Activase vial (29 million IU) is packaged with diluent for reconstitution (50 mL Sterile Water for Injection, USP): NDC 50242-044-13.

Each 100 mg Activase vial (58 million IU) is packaged with diluent for reconstitution (100 mL Sterile Water for Injection, USP), and one transfer device: NDC 50242-085-27.

### 16.2 Stability and Storage

Store lyophilized Activase at controlled room temperature not to exceed 30°C (86°F), or under refrigeration (2-8°C/36-46°F). Protect the lyophilized material during extended storage from excessive exposure to light. If stored between 2-30°C (36-86°F), Activase may be used within 8 hours following reconstitution. Discard any unused solution after administration is complete.

Do not use beyond the expiration date stamped on the vial.

## 17 PATIENT COUNSELING INFORMATION

Following Activase administration, patients are at increased risk of bleeding internally or externally. Advise patients to contact a health-care professional if they experience symptoms or signs consistent with bleeding (e.g., unusual bruising, pink or brown urine, red or black or tarry stools, coughing up blood, vomiting blood or blood that looks like coffee grounds), headache, or stroke symptoms.

### **Activase® (alteplase)**

Manufactured by:

**Genentech, Inc.**

A Member of the Roche Group

1 DNA Way

South San Francisco, CA

94080-4990

U.S. License No. 1048

Activase® is a registered trademark of Genentech, Inc.

©2018 Genentech, Inc.